Document Identifier: **PMC8144667**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: 2.2 Ethical statement The study was approved by the ethics committee of First Affiliated Hospital of Guangzhou University of Chinese Medicine (NO. ZYYECK 2017071).
Consent: Written informed consent was obtained from each participant.
Inclusion and Exclusion Criteria
Patients were excluded if there were neonatal anomalies, genital malformations, or uterine abnormalities (such as uterine fibroids or adenomyosis), abnormal amniotic fluid, had received drugs to inhibit uterine contractions within 48 h before cesarean section, postpartum hemorrhage, or other high-risk complications.
Attrition
not detected.
Sex as a biological variable
After explaining the aim of the study to eligible women face-to-face, the co-first author (JCZ) obtained written informed consent from those who had agreed to participate.
Subject Demographics
Weight: Patients were included in the study if they met the following criteria: cesarean section primipara, age >= 20 years and <= 40 years, single pregnancy, gave birth at >= 37 weeks of gestation, neonatal weight >= 2,500 g and < 4,000 g, no obstetric complications from pregnancy through the intrapartum period, no medical or surgical disease history, and could breastfeed after birth.
Age: AA was provided by a senior physician (SXW) with 18 years of clinical acupuncture experience.
Randomization
2.5 Randomization Patients from the postpartum ward at First Affiliated Hospital of Guangzhou University of Chinese Medicine were randomly allocated into either the real AA or sham AA groups in a 1:1 ratio.
Blinding
The researchers also performed blind evaluations of the patients after each treatment.
Power Analysis

2.4 Sample size calculation The sample size was calculated based on the research teams previous findings.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
PASS 11.0 software analyzed the experimental and control groups by comparing 2 independent means of the sample.	PASS		Suggestion: PASS (RRID:SCR_005490)(link)
All statistical analyses were performed using SPSS 20.0 (IBM SPSS Inc.,	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Group differences at each measuring time point were examined using 2 independent sample t-tests.	t-tests
The repeated measurements analysis of variance was used for comparison before and after treatment.	analysis of variance
Categorical variables, including categorical baseline variables and incidence of adverse events, were analyzed using the chi-square (kh2) test or the Fishers exact test.	chi-square
	Fishers exact test
Grade data was expressed as a composition ratio, and the 2 groups were compared using a Wilcoxon rank sum test.	Wilcoxon rank sum test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq