Document Identifier: **PMC8150440**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: The study was approved by the Independent Ethics Committee of the Bologna Health Authority (CE number: 09070).
Consent: Patients gave their written informed consent to participate in the study and to publish the data.
Inclusion and Exclusion Criteria
Secondary causes of parkinsonism were also excluded before enrollment by means of appropriate investigations including brain magnetic resonance imaging.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
All diagnoses were independently confirmed by three neurologists expert in movement disorders who were blinded to the diagnosis provided by the database.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
The main study outcome was the 3 h AUC_ETap calculated according to the linear trapezoidal rule by means of SigmaPlot 12.5 software (Systat Software, San Jose, CA, USA).	SigmaPlot		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical analysis When data were consistent with a normal distribution and equal variances, means and standard deviations were calculated and comparisons of study variables among patients subgroups were carried out by One Way Analysis of Variance (ANOVA).	Analysis of Variance
	ANOVA
Pairwise comparisons were performed by Holm-Sidak method when ANOVAs indicated a significant difference among subgroups.	Holm-Sidak
	ANOVAs
When deviation from a normal distribution was found, or when different var-iances were identified, medians and 25th-75th percentiles were calculated and comparisons were performed by Kruskal-Wallis One Way Analysis of Variance on Ranks (ANOVA on Ranks).	Kruskal-Wallis
Pairwise comparisons were performed by Dunns method when ANOVAs on Ranks indicated a significant difference among subgroups.	Dunns method
Sex and LIDs distribution was compared among groups by Chi square test.	Chi square test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq