Document Identifier: **PMC8153795**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: 2 Materials and Methods 2.1 Sample Collection and Protein Extraction The blood and resected ovarian biopsies were taken from 34 females diagnosed with OC and 20 with benign ovarian disease (Table S1, Section S1.1) after the approval of the University Ethical Review Board [ref No. 873/12].
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
2 Materials and Methods 2.1 Sample Collection and Protein Extraction The blood and resected ovarian biopsies were taken from 34 females diagnosed with OC and 20 with benign ovarian disease (Table S1, Section S1.1) after the approval of the University Ethical Review Board [ref No. 873/12].
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
All of the assays were performed in duplicate.

Number: All of the assays were performed in duplicate.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
2.4 Immunological Validation Western blotting (WB) was performed by fractionation of 20 µg of proteins by 12% SDS-PAGE followed by transfer to a nitrocellulose membrane, blocking with 5% BSA, and incubation with the primary rabbit polyclonal antibody to the tubulin beta 2C chain (1:500 dilution, PA5-25050, Invitrogen)	tubulin beta 2C chain		
	PA5-25050		
, vimentin (1:500 dilution , SAB 4503083 , Sigma-Aldrich) , and the anti-rabbit IgG horseradish peroxidase (HRP)-conjugated goat secondary antibody (1:6000 dilution , A6154 , Sigma-Aldrich)	vimentin		
	anti-rabbit IgG		
Software and Algorithms			
20 2.3 Gene Ontology Enrichment, Molecular Pathway, and Network Analysis The proteins exhibiting differential expression in OC were further categorized by gene ontology (GO) (www.geneontology.com) according to their molecular functions and biological processes and ingenuity pathway analysis (IPA, QIAGEN, www.qiagen.com/ingenuity) to identify molecular networks, canonical pathways, and the diseases and disorders associated with data set proteins.	ingenuity pathway analysis		Suggestion: Ingenuity Pathway Analysis (RRID:SCR_008653)(link)
Finally, IPA findings were validated by interacting protein network construction using STRING v10.0 (string.db.org) and Kyoto Encyclopedia of Genes and Genomes (KEGG) analysis.	STRING		Suggestion: STRING (RRID:SCR_005223)(link)
	KEGG		Suggestion: KEGG (RRID:SCR_012773)(link)
Protein bands were developed using a TMB (3,3#,5,5#-tetramethylbenzidine, Sigma) peroxidase substrate system.	Sigma		
2.5 Statistical Analysis Statistical analyses were done using SPSS version 15.0 (IBM Corporation, Armonk, NY) and/or GraphPad Prism 8.30.	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)
	GraphPad Prism		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
For selection of differentially stained 2D gel spots, the nonparametric Mann–Whitney U test was applied on summed-up intensities of the multiple spots of the same protein followed by false discovery rate (FDR) determination. 19 The proteins in 2D gel differential spots were digested in 20 ng/μL sequencing-grade trypsin (Promega, Southampton, UK).	Mann–Whitney U test
The Kruskal Wallace one-way analysis of variance (ANOVA) test was used for comparative analysis of the antibody titer among study cohorts with the significance level set at p-value < 0.05.	analysis of variance
	ANOVA

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq