Document Identifier: **PMC8168053**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: The present study was approved by the Ethics Committee of Sichuan Cancer Hospital (SCCHEC-02-2018-021), and all participants signed written informed consent forms.
Consent: The present study was approved by the Ethics Committee of Sichuan Cancer Hospital (SCCHEC-02-2018-021), and all participants signed written informed consent forms.
Inclusion and Exclusion Criteria
The following patients were excluded: (1) those treated with chemotherapy or radiotherapy prior to surgery, (2) those with prior history of other malignant tumors, and (3) those treated without radical surgery for tumor removal.
Attrition
not detected.
Sex as a biological variable
The following patients were excluded: (1) those treated with chemotherapy or radiotherapy prior to surgery, (2) those with prior history of other malignant tumors, and (3) those treated without radical surgery for tumor removal.
Subject Demographics
Age: Included patients in this study met the following criteria: (1) age <70 years, (2) diagnosed as pT3N0M0 ESCC, and (3) undergoing right thoracic radical resection.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication

not required.	
Data Information	
Identifiers: The quantity and quality of the extracted DNA were determined using a Qubit 3.0 Fluorometer (Thermo Fisher Scientific, Waltham, MA, USA, Q33217) and Agilent 2100 Bioanalyzer (Agilent Technologies, Palo Alto, CA, USA, G2939A), respectively.	Q33217

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Duplicate reads were marked by the Picard toolkit (version 2.1.0 , Welcome Trust Sanger Institute , Welcome Trust Genome Campus , Cambridge , CB10 1SA , UK) and then realigned using the Genome Analysis ToolKit (version 3.7 , Program in Medical and Population Genetics , Broad Institute of Harvard and MIT , Cambridge , MA , USA) .	Picard		
	Genome Analysis ToolKit		
, AstraZeneca , Waltham , MA , USA) , and the variants were annotated using ANNOVAR (Center for Applied Genomics , Childrens Hospital of Philadelphia , Philadelphia , PA , USA) .	ANNOVAR		Suggestion: ANNOVAR (RRID:SCR_012821)(link)
Compound heterozygous mutations were merged with FreeBayes (version 1.2.0 , Department of Biology , Boston College , Chestnut Hill , MA , USA) .	FreeBayes		Suggestion: FreeBayes (RRID:SCR_010761)(link)
Driver gene mutations and hotspot mutations included in the ExACCOSMIC database were filtered out .	ExACCOSMIC		
Reference genes and genome annotation files were available from the ENSEMBL website (http:www.ensembl.orgindex.html), and clean data were aligned to the reference genome by HISAT2 (http:ccb.jhu.edusoftwarehisat2index.shtml).	ENSEMBL		Suggestion: Ensembl (RRID:SCR_002344)(link)
	HISAT2		
HTSeq (http:www.huber.embl.deusersandersHTSeqdocoverview.html) was used to determine the expression level of each gene.	HTSeq		
Identification and functional enrichment analysis of differentially expressed genes (DEGs) Differential expression analysis of transcriptome sequencing data was performed using the DESeq2 package in R (www.r-project.org) .	DESeq2		Suggestion: DESeq, RRID:SCR_000154(link)
Selected DEGs were compared with Gene ontology and Kyoto Encyclopedia of Genes and Genomes databases, and the enrichment analysis of DEGs was conducted using Metascape (http:metascape.org).	Genomes		
	Metascape		

Statistical analysis The data were analyzed using R 3.5.1 and IBM SPSS Statistics for Windows, Version 22.0 (IBM Corp., Armonk, NY, USA).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
The Kaplan-Meier method and log-rank test were used for survival analysis and comparison, respectively.	log-rank test
The differences in SNV, CNV, TMB, MATH, and CNI between the different patient groups were analyzed by the Wilcoxon rank-sum test or Fishers exact test.	Wilcoxon rank-sum test
	Fishers exact test
Oligonucleotides	
The captured library was further amplified using Illumina p5 (5 AAT GAT ACG GCG ACC ACC GA 3) and p7 (5 CAA GCA GAA GAC GGC ATA CGA GAT 3) primers in the KAPA Hifi HotStart ReadyMix (KAPA Biosystems , KK2602) and purified with Agencourt AMPure XP beads .	5 AAT GAT ACG GCG ACC ACC GA 3
	CAA GCA GAA GAC GGC ATA CGA GAT 3

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq