Document Identifier: **PMC8170111**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not required.
Inclusion and Exclusion Criteria
not required.
Attrition
not required.
Sex as a biological variable
not detected.
Subject Demographics
Age: not required.
Weight: not required.
Randomization
Compound was added using the LabCyte Echo 555 acoustic dispenser in a randomised well layout at 4 hours post-seeding for G1max and 18 hours post-seeding for SG2max.
Blinding
not detected.
Power Analysis
not detected.
Replication
Each plate consisted of 12 DMSO control wells alongside an 11-point dose response (from 0.3 nM to 30 mM) for seven compounds with two replicates for each condition.

Type: Statistics Data is reported as mean with +- SEM or SD, detailed in the figure legend as applicable, of at least three independent biological replicates except for Figure 2 and 5 describing the compound library screen and scheduling comparison respectively where sample size is detailed in the figure legends.
Number: Each plate consisted of 12 DMSO control wells alongside an 11-point dose response (from 0.3 nM to 30 mM) for seven compounds with two replicates for each condition.
Cell Line Authentication
Contamination: Each cell line passed mycoplasma and single tandem repeat (STR) genotyping tests performed internally by the CRUK-CI Cell Services core facility.
Authentication: Each cell line passed mycoplasma and single tandem repeat (STR) genotyping tests performed internally by the CRUK-CI Cell Services core facility.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Fixed cells were then washed once with 1xPBS before being incubated in 500 mL blocking buffer (1xPBS +2 BSA + 0.1 TritonX-100Tween20) for 1 hour at room temperature (RT), primary antibody solution (phospho-MPM-2, 1:500, Merck Millipore 05-368 phospho-histone H2AX S139, 1:500, Merck Millipore 05-636) in blocking buffer for 2 hours at RT and secondary antibody solution (Alexa Fluor 488 Goat anti-mouse, 1:500, Thermo Fisher A11017) in blocking buffer for 1 hour at RT in the dark with three washes using blocking buffer between each step.	phospho-MPM-2		
	phospho-histone H2AX S139		
	anti-mouse		
Experimental Models: Cell Lines			
Cell culture & chemicals All cell lines were cultured at 37degC, 5 CO2 for a maximum of 20 passages after thawing and were cultured in DMEM (Thermo Fisher 41966029) (MIA PaCa-2, PANC-1, HeLa) or RPMI-1640 (Thermo Fisher 21875034) (AsPC-1) and supplemented with 10 fetal bovine serum (FBS) (Thermo Fisher 10270106).	PANC-1		Suggestion: (RRID:CVCL_A4BU)(link)
Proliferation assays For colony forming and clonocidal assays, 300 or 120 cells (MIA PaCa-2, HeLa, PANC-1, MCF7, K8484) and 1000 or 400 cells (AsPC-1) were seeded in 6 or 12 well plates respectively and grown for 11-14 days depending on the cell line with the final four days including drug exposure where indicated.	MCF7		Suggestion: RRID:CVCL_0031(link)
Repeat of a-d in three other cell lines, HeLa, MIA PaCa-2 and PANC-1. f.	HeLa		Suggestion: RRID:CVCL_0030(link)
Software and Algorithms			
RO-3306 (Merck, 217699), nocodazole (Sigma-Aldrich, M1404), gemcitabine (Tocris, 3259), CB-5083 (BioVision via Cambridge Bioscience, B1032-5), palbociclib (Sigma, PZ0199) and SN-38 (Merck,	Sigma		Suggestion: sigma-tau (RRID:SCR_000488)(link)

H0165) were dissolved in DMSO, aliquoted and stored at -20degC.			
Fixed cells were then incubated with blocking solution containing the DNA dye FxCycle Violet (1 mgmL, Thermo Fisher F10347) for at least 1 hour at RT in the dark before samples were run using the BD Biosciences LSRFortessa(tm) flow cytometer and analysed using FlowJo(r) and GraphPad Prism software.	GraphPad Prism		
PRISM 8.4.0 (GraphPad Software, San Diego, CA, USA) was used for data analysis.	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical analysis of two parameters was performed using either a two-tailed, unpaired t test (Figure 1d), or two-tailed, unpaired Mann-Whitney test (Figure 4de) based on data distribution.	t test
	Mann-Whitney test
For multiple comparisons involving more than two parameters, a multiple unpaired t test with Holm-Sidak multiple comparisons correction (Figure 1b) or a one-way or two-way ANOVA was performed with a Bonferroni correction (Figure 3b, 4df) were performed.	Holm-Sidak multiple comparisons correction
	ANOVA
	Bonferroni
Students t tests (c) or multiple t tests (b) with a Holm-SIdak multiple comparisons correction was performed for comparisons between two groups or more respectively.	t tests

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq