Document Identifier: **PMC8170869**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: Animal experiments were carried out at The First Affiliated Hospital of Chongqing Medical University (Chongqing, China) and were approved by the Animal Ethics and Use Committee of the First Affiliated Hospital of Chongqing Medical University (approval no. 2017-028).
Euthanasia Agents: Following anesthetization with 3% sodium pentobarbital (80 mg/kg), five mice from each group were decapitated to collect the blood samples after 24, 48 and 72 h of reperfusion.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
Materials and methods Animals and grouping A total of 45 male C57BL/6J mice (age, 8–12 weeks; weight, 20–25 g) were purchased from Vital River Laboratories.
Subject Demographics
Age: Materials and methods Animals and grouping A total of 45 male C57BL/6J mice (age, 8–12 weeks; weight, 20–25 g) were purchased from Vital River Laboratories.
Weight: Materials and methods Animals and grouping A total of 45 male C57BL/6J mice (age, 8–12 weeks; weight, 20–25 g) were purchased from Vital River Laboratories.
Randomization
The mice were randomly divided into the sham-surgery, IRI, CXCR3 overexpression (OE-CXCR3)+IRI, PC61+IRI and OE-CXCR3+PC61+IRI groups (n=15 mice/group).
Blinding
not detected.
Power Analysis
not detected.

Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
In brief, the aforementioned cells isolated from the kidneys were incubated with the anti-CD4 antibody (cat. no. 11-0041-82; FITC; eBioscience; Thermo Fisher Scientific, Inc.) at a dilution of 1:10 for 1 h at 4°C, followed by incubation with Cytofix/Cytoperm (BD Biosciences) at 4°C for 15 min to permeabilize the cell membranes.	anti-CD4		Suggestion: (DSHB Cat# anti-CD38 (HB-7), RRID:AB_3105832)(link)
The cells were then stained with PE-conjugated anti-Foxp3 antibodies (cat. no. 12-5773-82; eBioscience; Thermo Fisher Scientific, Inc.) at 1:20 dilution for 1 h at 4°C and analyzed on a BD FACSCalibur System (BD Bioscience).	anti-Foxp3		Suggestion: (DSHB Cat# PCRP-FOXP3-1B2, RRID:AB_2618638)(link)
The PVDF membranes were incubated with 5% fat-free milk for 1 h at room temperature and then incubated with the primary antibodies overnight at 4°C, including anti-fibronectin antibody (1:2,000 dilution; cat. no. ab2413; Abcam), anti-type IV collagen antibody (1:2,000 dilution; cat. no. ab19808; Abcam), anti-IL-6 antibody (1:2,000 dilution; cat. no. sc-32296; Santa Cruz Biotechnology, Inc.), anti-TNF-α antibody (1:2,000 dilution; ab183218; Abcam), anti-CCL-2 antibody (1:2,000 dilution; cat. no. 2029; Cell Signaling Technology, Inc.), anti-IL-10 antibody (1:4,000 dilution; ab189392; Abcam), anti-Bax antibody (1:3,000 dilution; cat. no. ab32503; Abcam), anti-Bcl-2 antibody (1:3,000 dilution; cat. no. ab182858; Abcam), anti-cleaved caspase-3 antibody (1:3,000 dilution; cat. no. ab49822; Abcam) and anti-β-actin antibody (1:5,000 dilution; cat. no. AM4302; Thermo Fisher Scientific, Inc.).	anti-fibronectin		
	anti-type IV collagen		
	anti-IL-6		
	anti-TNF-α		
	anti-CCL-2		
	anti-IL-10		
	anti-Bax		
	anti-Bcl-2		
	anti-cleaved caspase-3		
	anti-β-actin		
Experimental Models: Organisms/Strains			
Materials and methods Animals and grouping A total of 45 male C57BL/6J mice (age, 8–12 weeks;	C57BL/6J		Suggestion: RRID:IMSR_JAX:000664(link)

weight, 20–25 g) were purchased from Vital River Laboratories.			
Preparation of the single-cell suspensions from the kidneys Single-cell suspensions of kidney cells were isolated from C57BL/6 mice based on a previous study (19).	C57BL/6		Suggestion: RRID:IMSR_JAX:000664(link)
Software and Algorithms			
The cells were then stained with PE-conjugated anti-Foxp3 antibodies (cat. no. 12-5773-82; eBioscience; Thermo Fisher Scientific, Inc.) at 1:20 dilution for 1 h at 4°C and analyzed on a BD FACSCalibur System (BD Bioscience).	BD FACSCalibur System		
Tregs were considered as CD4+ and Foxp3+ and was analyzed using the FlowJo 7.6 software (FlowJo LLC).	FlowJo		Suggestion: FlowJo, RRID:SCR_008520(link)
The gray-scale value analysis was performed using ImageJ software (version 1.48; National Institutes of Health).	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
Statistical analyses were performed using SPSS software (version 21.0; IBM Corp.).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Two-sided unpaired t-tests and one-way ANOVA followed by Bonferroni post-hoc tests were used to compare differences between 2 and ≥3 groups, respectively.	t-tests
	ANOVA
	Bonferroni post-hoc tests
Kidney injury score was expressed as median and range, and was analyzed using the Kruskal-Wallis test followed by the Steel-Dwass test.	Kruskal-Wallis test
	Steel-Dwass test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq