Document Identifier: **PMC8178775**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not required.
Inclusion and Exclusion Criteria
not required.
Attrition
GSE93249 included 11 rat spinal cord samples, including 3 non-injured spinal cord samples in the sham group and 8 SCI samples in the SCI group, representing the chronic phase at 1 month (m1, n = 3), 3 months (m3, n = 3), and 6 months (m6, n = 2) after injury.
Sex as a biological variable
not required.
Subject Demographics
Age: not required.
Weight: not required.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.
Data Information

Identifiers: Materials and Methods Microarray data preprocessing The gene expression profiles GSE45006, GSE45550, and GSE93249 were downloaded from the Gene Expression Omnibus database of the National Center of Biotechnology Information (http:www.ncbi.nlm.nih.govgeo).	GSE45006
	GSE45550
	GSE93249

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Materials and Methods Microarray data preprocessing The gene expression profiles GSE45006, GSE45550, and GSE93249 were downloaded from the Gene Expression Omnibus database of the National Center of Biotechnology Information (http:www.ncbi.nlm.nih.govgeo).	Gene Expression Omnibus		Suggestion: Gene Expression Omnibus (GEO) (RRID:SCR_005012)(link)
The identification of DEGs between SCIs and sham samples The DEGs between SCI and sham samples were screened using GEO2R (http:www.ncbi.nlm.nih.govgeogeo2r) online tools (Davis and Meltzer, 2007).	GEO2R		Suggestion: GEO2R (RRID:SCR_016569)(link)
GO and KEGG pathway enrichment analyses of the DEGs GO analysis has been widely used to identify functional enrichment patterns in studies examining large-scale genetic sequencing.	KEGG		
The Database for Annotation, Visualization, and Integrated Discovery (DAVID) provides a comprehensive set of functional annotation tools.	DAVID		
Statistical analysis SPSS 22.0 (IBM, Armonk, NY, USA) was used for all statistical analyses.	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
The false discovery rate (FDR) was obtained by adjusting the raw P-values with the Benjamini-Hochberg method (Madar and Batista, 2016).	Benjamini-Hochberg
Comparisons among multiple groups were analyzed using a one-way analysis of variance, followed by Dunnetts post hoc test.	analysis of variance
	Dunnetts post hoc test
For experiments involving two groups, an unpaired Students t-test was performed.	t-test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq