Document Identifier: **PMC8180462**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: The experimental protocol was approved by the institutional of Animals Care and Use Committee (ACUC) and Research Ethics Committee (REC), reference number-595-17, Faculty of medicine, King Abdulaziz University, Jeddah, Saudi Arabia.
IRB: The experimental protocol was approved by the institutional of Animals Care and Use Committee (ACUC) and Research Ethics Committee (REC), reference number-595-17, Faculty of medicine, King Abdulaziz University, Jeddah, Saudi Arabia.
Euthanasia Agents: Subsequently, blood samples were collected from the ophthalmic venous plexus through retro-orbital approach, and the animals were decapitated.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
2.1 Study design Fifty male Wistar rats, 6–8 weeks old with 180–250 g body weight, were chosen for the study.
Subject Demographics
Age: 2.1 Study design Fifty male Wistar rats, 6–8 weeks old with 180–250 g body weight, were chosen for the study.
Weight: 2.1 Study design Fifty male Wistar rats, 6–8 weeks old with 180–250 g body weight, were chosen for the study.
Randomization
Rats were randomly divided into five groups: Group A:(control-negative) received vehicle for VPA and ASTA by oral gavage daily and vehicle for PTZ by intraperitoneally three times a week, Group B: (control-positive) received vehicle for VPA and ASTA by oral gavage daily and 30 mg/kg of PTZ (Atack et al., 2000) by intraperitoneally three times a week, Group C: received 500 mg/kg of valproic acid (El-Mowafy et al., 2016) by oral gavage daily and PTZ by intraperitoneally three times a week, Group D: received 100 mg/kg of astaxanthin (Chang et al., 2018) by oral gavage daily and, PTZ by intraperitoneally three times a week, and

Group E: received valproic acid, astaxanthin by oral gavage daily 2 h apart and PTZ by intraperitoneally three times a week.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Organisms/Strains			
2.1 Study design Fifty male Wistar rats, 6–8 weeks old with 180–250 g body weight, were chosen for the study.	Wistar		Suggestion: RRID:RGD_13508588(link)
Software and Algorithms			
2 Material and methods Valproic acid was obtained from Sigma Aldrich (CAS number: 1069-66-5), while Astaxanthin (Astaxanthin-Enriched Haematococcus pluvialis powder 5% (AstaZine)) was a gift from Algea health sciences INC; A BGG company (CAS number 472-61-7).	Sigma Aldrich		Suggestion: Sigma-Aldrich (RRID:SCR_008988)(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical analysis was performed using one-way analysis of variance (ANOVA) with Tukey's post hoc test.	analysis of variance
	ANOVA
	Tukey's post hoc test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq