Document Identifier: **PMC8203849**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Ethics statement The Institutional Review Boards of the Korean National Institute of Health for the Korean Genome and Epidemiology Study (No.: KBP-2015-055) and Hoseo University (No.: 1041231-150811-HR-034-01) approved the study.
Consent: All that participated in KoGES provided written informed consent.
Inclusion and Exclusion Criteria
A variable to be analyzed was excluded from the covariates.
Attrition
not detected.
Sex as a biological variable
Definition of MS MS was defined as the presence of a cluster of >= 3 of the following: 1) abdominal obesity (waist circumference >= 90 cm for men and >= 85 cm for women) 2) elevated blood pressure (average systolic blood pressure >= 130 mmHg or diastolic blood pressure >= 85 mmHg) 3) low HDL-cholesterol level (< 40 mgdL for men and < 50 mgdL for women) 4) elevated serum triglyceride level (>= 150 mgdL) and 5) elevated fasting blood glucose level (>= 110 mgdL).
Subject Demographics
Age: METHODS Participant characteristics and biochemical results A total of 40,262 Korean adults aged 51-77 years participated from 2004-2013 in the KoGES, organized by the Korean Center for Disease and Control.13 General characteristics, including socioeconomic factors, were acquired using survey
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.

Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
17 18 Daily intakes of energy and nutrients (e.g. , protein , carbohydrate , and fat) were calculated using Can-Pro 2.0 nutrient assessment software developed by the Korean Nutrition Society.17 The 106 food items in SQFFQ were categorized into 30 predefined food groups (Supplementary Table 1) , and principal component analysis (PCA) was conducted with food groups using proc principal component and proc factor procedure.	Can-Pro		
Statistical analysis Statistical analysis was performed using SAS version 9.4 (SAS Institute, Cary, NC, USA).	SAS Institute		Suggestion: Statistical Analysis System, RRID:SCR_008567(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
According to MS status, frequency distributions of classification variables were calculated, and statistical analysis was conducted using the chi-squared test.	chi-squared test
Means and standard errors of general characteristics were calculated, and their statistical differences were analyzed with two-way analysis of variance with main effects of ARC and MS.	analysis of variance
According to MS and ARC, the significance of MS components was determined by a two-way analysis of covariance (ANCOVA) with covariates adjustments.	analysis of covariance
	ANCOVA
Intergroup multiple comparisons were performed using the Tukey test.	Tukey test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq