Document Identifier: **PMC8232843**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: Methods Ethics Statement Animal experiments were authorized by the Committee of Experimental Animals of Hengdian Wenrongs Hospital (Z20190921H).
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
Mice Fifty female BALBc mice (weight: 20-25 g, age: 6-8 weeks) were obtained from SLAC Laboratory Animal Technology (Shanghai, China).
Subject Demographics
Weight: Mice Fifty female BALBc mice (weight: 20-25 g, age: 6-8 weeks) were obtained from SLAC Laboratory Animal Technology (Shanghai, China).
Age: Mice Fifty female BALBc mice (weight: 20-25 g, age: 6-8 weeks) were obtained from SLAC Laboratory Animal Technology (Shanghai, China).
Randomization
Treatment of the Experimental Mice The experimental mice were randomly divided into the following 5 groups (n=10): the Sham group, the Curcumol-H group, the OVA group, the OVA+Curcumol-L group, and the OVA+Curcumol-H group.
Blinding
not detected.
Power Analysis
not detected.
Replication

not required.	
Data Information	
Identifiers: Then, the protein swam vertically down inside the gels under 100 volt for 120 min, and then transferred to the surface of 0.45 mm membranes (ISEQ00010, Millipore, MA, USA).	ISEQ00010

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
The second day, the slices were further incubated with goat-anti-rabbit IgG (1:30000, ab205718, Abcam) antibody for 1 min, followed by incubation with the DBA reagent (SFQ004, 4A Biotech, Beijing, China) for 30 min.	IgG		Suggestion: (Cytognos Cat# CYT-IGG4AP, RRID:AB_2876879)(link)
	SFQ004		
The relative primary antibodies were Wnt5a (1:1500, 42kD, ab229200, Abcam), b-catenin (1:1000, 86kD, ab32572, Abcam), VEGFA (1:2500, 27kD, ab52917, Abcam), TGF-b1 (1:1000, 44kD, ab92486, Abcam), Fibronectin (1:1500, 262kD, ab2413, Abcam), MMP-9 (1:2500, 92kD, ab38898, Abcam), and GAPDH (1:5000, 36kD, ab181602, Abcam).	Wnt5a		
	b-catenin		
	VEGFA		
	TGF-b1		
	Fibronectin		
	MMP-9		
	GAPDH		
Software and Algorithms			
Statistical Analysis All the statistical analyses were conducted using GraphPad 8.0 software.	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)
Q-Q plot was performed by IBM SPSS Statistics 20.	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Comparison between multiple groups was analyzed using One-way ANOVA.	ANOVA
Shapiro-Wilk test was used to assess data distribution.	Shapiro-Wilk test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq