Document Identifier: **PMC8236216**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Isolation and culture of human primary T cells Primary adult blood cells were obtained from anonymous healthy human donors as a leukoreduction pack purchased from StemCell Technologies , Inc . or Allcells Inc , or as a Trima residual from Vitalant , under a protocol approved by the University of California , San Francisco Institutional Review Board (IRB) .
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.
Cell Line Authentication

Contamination: Cell lines were routinely checked for mycoplasma using the MycoAlert mycoplasma detection kit (Lonza) according to the manufacturer’s instructions .	
Data Information	
Availability: Materials availability Plasmids generated in this study have been deposited to Addgene or are available upon request.	
Identifiers: PRJNA680046) .	PRJNA680046
Identifiers: HIV-1 Env amino acid sequence was obtained from UniProt (P04578) , human codon optimized (IDT) , and ordered as a gBlock (IDT) .	P04578

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
96-well flat bottom plates were primed for stimulation by incubating with anti-human CD3 (10 µg/mL) and anti-human CD28 (5 µg/mL) antibodies in PBS for 1 hour at 37°C prior to washing .	anti-human CD3		
	anti-human CD28		
The solution was replaced with 1 % non-fat milk in PBS-T and a 1:5000 primary antibody dilution containing anti-FLAG (Sigma) or a 1:2000 dilution of anti-p24 (Abcam) antibodies prior to shaking at 4°C overnight .	anti-FLAG		
	anti-p24 (Abcam)		
Cells were resuspended in FACS buffer (1 % –2 % BSA in PBS) and stained with the surface marker-targeting antibodies: B2M-FITC (Biolegend) , B2M-PE (Biolegend) , B2M-APC (Biolegend) , CD4-FITC (Biolegend) , CD8-PeCy7 (BD Biosciences) , and TCRa/b-BV421 (Biolegend) and live/dead stains Ghost Dye red 780 (Tonbo) or Ghost Dye violet 450 (Tonbo) , prior to analyzing .	B2M-PE		
	B2M-APC		
	CD4-FITC		
	CD8-PeCy7		
	TCRa/b-BV421		
CD69 PerCP (BioLegend) , 4-1BB BV711 (BioLegend) and intracellular antibodies TNF-a Pacific Blue (BioLegend) , IL-2 APC (BD) and IFN-g FITC (BioLegend) using the FIX & PERM Cell Fixation &	BV711		
	IL-2 APC		
	IFN-g FITC		
Experimental Models: Cell Lines			
EXPERIMENTAL MODEL AND SUBJECT DETAILS Culture of human cell lines Lenti-X , 293T , A549 , CCRF-CEM , HuT 78 and Jurkat cell lines were obtained from the UC Berkeley Cell Culture Facility .	Jurkat		
Lenti-X , 293T , and A549 cells were cultured in DMEM (Corning) , Jurkat and CCRF-CEM cells were cultured in RPMI 1640 (Thermo Fisher) and 1 mM sodium pyruvate , while HuT 78 cells were cultured in IMEM (Thermo Fisher)	A549		Suggestion: RRID:CVCL_0023(link)
	HuT 78		Suggestion: (RRID:CVCL_0337)(link)
Targeted integration analysis 15k 293T cells treated with B2M-targeting or non-targeting Cas9-VLPs	293T		Suggestion: RRID:CVCL_0063(link)

and DNA was isolated 3 days post treatment by resuspending in Quick Extract (Lucigen) and heating at 65°C for 20 minutes followed by 95°C for 20 minutes before storing at –20°C .			
Recombinant DNA			
METHOD DETAILS Plasmid construction The Gag-pol expression plasmid psPax2 was a gift from Didier Trono (Addgene plasmid #12260) .	psPax2		
pCMV-VSV-G was a gift from Bob Weinberg (Addgene plasmid #8454) .	pCMV-VSV-G		
Gag-Cas9 was constructed by amplifying Gag from psPax2 and Cas9 from pMJ920 (	pMJ920		
In-Fusion (Takara Bio) cloning was used to clone Gag-Cas9 and Env into the pCAGGS expression vector .	pCAGGS		
pCF221 (Addgene plasmid #121669) was modified to express mNeonGreen (Allele Biotechnology) or the #-CD19-4-1BBζ-P2A-mCherry (CAR-P2A-mCherry) construct (Hill et al. , 2018; Muller et al. , 2020) in place of mCherry and was used as the sgRNA-expressing lentiviral transfer plasmid.	pCF221		
For generation of hybrid Cas9-VLPs , the guide RNA expression cassette was removed from the CAR-P2A-mCherry lentiviral plasmid via digestion with EcoRI and KpnI (NEB) .	CAR-P2A-mCherry		
The U6-sgRNA CAG-mTagBFP2 expression plasmid used for traceless Cas9-VLP and CAR-Cas9 VLP production was a gift from Benjamin Oakes	CAG-mTagBFP2		
Software and Algorithms			
All cells were cultured with 10 % fetal bovine sera (VWR) and 100 U/mL penicillin-streptomycin (GIBCO) .	GIBCO		
HIV-1 Env amino acid sequence was obtained from UniProt (P04578) , human codon optimized (IDT) , and ordered as a gBlock (IDT) .	UniProt		
The solution was replaced with 1 % non-fat milk in PBS-T and a 1:5000 primary antibody dilution	Sigma		

containing anti-FLAG (Sigma) or a 1:2000 dilution of anti-p24 (Abcam) antibodies prior to shaking at 4°C overnight .			
All analysis was done using the FlowJo v10 software.	FlowJo		Suggestion: FlowJo, RRID:SCR_008520(link)
QUANTIFICATION AND STATISTICAL ANALYSIS Statistical analysis Statistical analysis was performed in Prism v7,v8, and v9.	Prism		Suggestion: Meta-Prism (RRID:SCR_021836)(link)

Other Entities Detected

Your Sentences	Recognized Entity
Unresolved RRIDs	
METHOD DETAILS Plasmid construction The Gag-pol expression plasmid psPax2 was a gift from Didier Trono (Addgene plasmid #12260) .	Addgene plasmid #12260
pCMV-VSV-G was a gift from Bob Weinberg (Addgene plasmid #8454) .	Addgene plasmid #8454
Addgene plasmid #42234) .	Addgene plasmid #42234
pCF221 (Addgene plasmid #121669) was modified to express mNeonGreen (Allele Biotechnology) or the #-CD19-4-1BBζ-P2A-mCherry (CAR-P2A-mCherry) construct (Hill et al. , 2018; Muller et al. , 2020) in place of mCherry and was used as the sgRNA-expressing lentiviral transfer plasmid.	Addgene plasmid #121669
The mixture was electroporated using a 4D-nucleofector (Lonza) before immediately adding to 293T cells stably expressing a BFP-to-GFP reporter (Addgene plasmid #71825) .	Addgene plasmid #71825

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq