Document Identifier: **PMC8236919**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not required.
Inclusion and Exclusion Criteria
not required.
Attrition
not required.
Sex as a biological variable
not detected.
Subject Demographics
Age: not required.
Weight: not required.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
Number: Statistical analysis Statistical results were obtained from at least three independent experiments.
Cell Line Authentication
not detected.

Code Information	
Availability: Plasmids are listed in Table S5, and SnapGene files will be provided upon request.	
Identifiers: Ridgeline plots were drawn in RStudio with ggplot2 and ggridges (https://github.com/clauswilke/ggridges) in R 3.5.3 .	https://github.com/clauswilke/ggridges

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Cell Lines			
These vectors are packaged by a third-generation system comprising four distinct packaging vectors (Tat , Rev , Gag/Pol , and either vesicular stomatitis virus G or murine leukemia virus envelope) using HEK 293T cells as the host .	HEK 293T		Suggestion: RRID:CVCL_0063(link)
DNA (plasmid of interest , 5 µg; Tat , 0.5 µg; Rev , 0.5 µg; Gag/Pol , 0.5 µg; vesicular stomatitis virus G or murine leukemia virus envelope , 1 µg) was delivered to the HEK cells as calcium phosphate precipitates .	HEK		Suggestion: (KCB Cat# KCB 86020YJ, RRID:CVCL_WZ56)(link)
Recombinant DNA			
Materials and methods Plasmids Plasmids were assembled by Gibson assembly (New England Biolabs) or the T5 exonuclease DNA assembly method (Xia et al., 2019) into the pHAGE lentiviral backbone (Wilson et al., 2008).	pHAGE		Suggestion: RRID:Addgene_166468(link)
Lentivirus production Lentiviral vector–packaged pHAGE-derived plasmids (Wilson et al. , 2008) were used for transfection .	pHAGE-derived		
Software and Algorithms			
All inserts were fully sequenced and matched to the Ensembl reference sequence, National Center for Biotechnology Information reference sequence, or expected mutant forms.	Ensembl		
Plasmids are listed in Table S5, and SnapGene files will be provided upon request.	SnapGene		
These cells were cultured in 95% DMEM (4.5 g/liter glucose), 5% FBS, 100 U/ml penicillin, and 100 µg/ml streptomycin (all from Gibco or Invitrogen).	Gibco		
MA; Addgene plasmid 52961; Sanjana et al. , 2014) or lentiCRISPR v2 PuroP93S (BL245) and screened by sequencing . lentiCRISPR v2 PuroP93S is similar to its parent except for a proline-to-serine mutation in the Puro N-acetyltransferase gene , which	https:// www.addgene.org		

increases its resistance to Puro (https://www.addgene.org).			
The template for homology-directed repair was designed in Benchling (https://www.benchling.com/).	Benchling		
Data were processed in FlowJo version 10 .	FlowJo		Suggestion: FlowJo, RRID:SCR_008520(link)
Ridgeline plots were drawn in RStudio with ggplot2 and ggridges (https://github.com/clauswilke/ggridges) in R 3.5.3 .	RStudio		
	ggplot2		Suggestion: ggplot2 (RRID:SCR_014601)(link)
Three-image Z stacks were acquired at 0.2-micron intervals and converted to single planes by maximum projection with MetaMorph software (MDS Analytical Technologies) .	MetaMorph		
Images were processed in ZEN 3.1 blue edition (Zeiss) or Adobe Photoshop CS6 with brightness and contrast adjusted.	ZEN		
	Adobe Photoshop		Suggestion: Adobe Photoshop, RRID:SCR_014199(link)
Statistical differences between groups were tested by one-way ANOVA, two-way ANOVA, or repeated measures ANOVA in GraphPad Prism 7.04.	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical differences between groups were tested by one-way ANOVA, two-way ANOVA, or repeated measures ANOVA in GraphPad Prism 7.04.	ANOVA
Oligonucleotides	
The following primers were used for detecting Ty1 knock-in of Ptch1: Ptch1KI_1 , 5#-TGTGTTGCTCCGAGC TTTGT-3#; Ptch1KI_2 , 5#-TCCATCCGG ACTCCAGACAT-3#; Ptch1KI_3 , 5#-GAACTTCAGCATCAAGCGGAT-3#; Ptch1KI_4 , 5#-CATTTGCTTCTGGCC ACACC-3#; Ptch1KI_5 , 5#-AAAAGG TCCCCCAGGGTCTA-3#; Ptch1KI_6 , 5#-ATGACCTCAGAAGCTGCCAC-3# .	5#-TGTGTTGCTCCGAGCTTTGT-3#
	5#-TCCATCCGGACTCCAGACAT-3#
	5#-GAACTTCAGCATCAAGCGGAT-3#
	5#-CATTTGCTTCTGGCCACACC-3#
	5#-AAAAGGTCCCCCAGGGTCTA-3#
	5#-ATGACCTCAGAAGCTGCCAC-3#
Unresolved RRIDs	
MA; Addgene plasmid 52961; Sanjana et al. , 2014) or lentiCRISPR v2 PuroP93S (BL245) and screened by sequencing . lentiCRISPR v2 PuroP93S is similar to its parent except for a proline-to-serine mutation in the Puro N-acetyltransferase gene , which increases its resistance to Puro (https://www.addgene.org).	plasmid 52961

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq