Document Identifier: **PMC8246653**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not required.
Inclusion and Exclusion Criteria
Inclusion criteria are RCTs conducted on T2DM patients with NAFLD on treatment with antidiabetic drugs, namely SGLT2Is, along with its effects on NAFLDNASH.
Attrition
not required.
Sex as a biological variable
not required.
Subject Demographics
Age: not required.
Weight: not required.
Randomization
75 years) Multi Double-blind 2 12 weeks 12 weeks Eriksson et al, 18 2018 Sweden Dapagliflozin 10 mg Placebo Patients with T2DM Multi Double-blind 2 12 weeks 12 weeks Guja et al, 23 2018 USA Dapagliflozin Exenatide Adults with T2DM and inadequateglycemic control Multi Double-blind 3 28 weeks 28 weeks Hayashi et al, 24 2017 Japan Dapagliflozin 5 mg Sitagliptin 50 mg Patients with T2DM taking prescribed oral hypoglycemic agents Single Open-label 2 12 weeks 12 weeks Kurinami et al, 38 2018 Japan Dapagliflozin 5 mg Non-SGLT2 (din specify) Patients with T2DM Single Open-lable 2 24 weeks 24 weeks Leiter et al, 27 2016 Germany Canagliflozin 300 mg Sitagliptin 100 mg Patients with T2DM Multi Double-blind 3 26 weeks 26 weeks Leiter et al, 28 2016 Germany Canagliflozin 100 mg Placebo Patients with T2DM Multi Double-blind 3 26 weeks 26 weeks Leiter et al, 28 2016 Germany Canagliflozin 300 mg Placebo Patients with T2DM Multi Double-blind 3 26 weeks 26 weeks Polidori et al, 29 2017 USA Canagliflozin 100 mg Placebo Patients with T2DM Single Unknown 3 26 weeks 26 weeks Polidori et al, 29 2017 USA Canagliflozin 300 mg Placebo Patients with T2DM Single Unknown 3 26 weeks 26 weeks Seko et al, 7 2017 Japan Canagliflozin 100 mg Placebo Patients with T2DMwith Baseline ALT ?? 30 Multi Double-blind 3 12 weeks 12 weeks Seko et al, 7 2017 Japan Canagliflozin 100 mg Placebo Patients with T2DMBaseline ALT > 30 Multi Double-blind 3 12 weeks 12 weeks Shimizu et al, 30 2018 Japan Dapagliflozin 5 mg Control (did not specify) Patients with T2DMand NAFLD Single Open-label 2 24 weeks 24 weeks RCT: randomized controlled trial T2DM: type 2 diabetes mellitus non-alcoholic fatty liver disease.

Blinding
not detected.
Power Analysis
not detected.
Replication
Number: Three independent authors preliminarily screened articles by their titles and abstracts, followed by full-text reading.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
A systematic literature search was performed using three databases, Medline, Cochrane, and Embase, from inception to 20 October 2018.	Medline		Suggestion: MEDLINE (RRID:SCR_002185)(link)
	Cochrane		Suggestion: Cochrane Library (RRID:SCR_013000)(link)
	Embase		Suggestion: EMBASE (RRID:SCR_001650)(link)

Other Entities Detected

Your Sentences	Recognized Entity
Oligonucleotides	
The primary outcome for these meta-analyses was the changes in hepatic enzymes levels, namely ALT, AST, and GGT.	GGT

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq