Document Identifier: **PMC8247603**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: All mice were maintained according to UCSF IACUC standards.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
MMFs were isolated from 6- to 10-wk-old female C57B6 Atg5flfllsl-RFP++ , Atg12flfllsl-RFP++ , or lsl-RFP ++ mouse mammary glands as previously described (Proia and Kuperwasser 2006).
Subject Demographics
Age: MMFs were isolated from 6- to 10-wk-old female C57B6 Atg5flfllsl-RFP++ , Atg12flfllsl-RFP++ , or lsl-RFP++ mouse mammary glands as previously described (Proia and Kuperwasser 2006).
Randomization
not detected.
Blinding
For tumor studies, mice were monitored for mammary tumor development, and blind measurements were obtained using calipers.
Power Analysis
not detected.
Replication
A ratio of 200 uL of diluted , neutralized collagen was used per 100 uL of MMF cell suspension for each technical replicate 300 uL of diluted , neutralized collagenMMF mixture was seeded into one well of a 24-well plate.

Type: A ratio of 200 uL of diluted , neutralized collagen was used per 100 uL of MMF cell suspension for each technical replicate 300 uL of diluted , neutralized collagenMMF mixture was seeded into one well of a 24-well plate.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Membranes were blocked for 1 h in 5 milk in PBS with 0.1 Tween 20 (PBST), incubated with primary antibody in 5 milkPBST overnight at 4degC, washed, incubated for 1 h at room temperature with HRP-conjugated donkey secondary antibodies (Jackson ImmunoResearch) at 1:5000 in 5 milkPBST, washed, and developed with Luminol signal reagent (Fisher).	HRP-conjugated donkey secondary		
Primary antibodies used for immunoblotting were as follows: ATG12 (1:500 Cell Signaling 2011), ATG5 (1:2000 Novus Biologicals NB110-53818)	ATG12		
	ATG5		
Primary antibodies used for immunohistochemistry were as follows: phosphorylated histone H3 (pH 3 1:100 Cell Signaling 9701) , p62 (1:200 Progen Biotechnik GP62-C) ,	phosphorylated histone H3 (pH 3 1:100 Cell Signaling 9701) , p62		
Primary antibodies used included aSMA (1:200 Vector Laboratories VP-S281) and S100A4 (1:50 Abcam 27957) .	S100A4		
Cells were stained using anti-LAMP1 (1:2000 BD Pharmingen 1D4B) and type 1 procollagen (PC1 1:100 LF42) overnight at 4degC, and Alexa Fluor secondary antibodies were used (1:1000 and 1:500, respectively).	anti-LAMP1		
	PC1		
Experimental Models: Organisms/Strains			
Animal models The generation of Atg12flfl mice has been described (Malhotra et al. 2015).	Atg12flfl		
Atg5flfl mice were obtained from Noboru Mizushima (Kuma et al. 2004).	Atg5flfl		
R26-lsl-tdRFP mice were obtained from Hans Joerg Fehling (Luche et al. 2007), bred to Atg12flfl and Atg5flfl mice, and backcrossed for more than five generations to C57B6 mice.	R26-lsl-tdRFP		
FSP-Cre (FvB) and MMTV-PyMT (C57B6) mice were provided by Zena Werb (UCSF).	C57B6		Suggestion: RRID:IMSR_JAX:000664(link)
FSP-Cre mice were bred to the Atg12flfl and Atg5flfl strains and	FSP-Cre		

then bred to MMTV-PyMT males to obtain Atg12flfllsl-RFP++PyMT +FSP-Cre+ and Atg5flfllsl-RFP+ +PyMT+FSP-Cre+ mice.	MMTV-PyMT		
	Atg5flfllsl-RFP+ +PyMT+FSP-Cre +		
In vivo MFP priming assays C57Bl6 mice (22 d old) were purchased from Jackson Laboratories (strain no. 000664).	C57Bl6		
Software and Algorithms			
Densitometry was performed with ImageJFiji software.	ImageJFiji		
Collagen was quantified using a set threshold in ImageJ and measuring more than four regions per primed MFP. H&E and Massons trichrome staining These stains were performed by the University of California at San Francisco Mouse Pathology Core and scanned with an Aperio ScanScope XT slide scanner (Aperio Technologies) at 40x.	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
The force data were converted to elastic modulus using the Hertz Model program (tissue samples assumed to be incompressible and a Poissons ratio of 0.5 was used to calculate Youngs elastic modulus) in IgorPro v6.22, supplied by Asylum Research.	IgorPro		
Bafilomycin or DMSO (Sigma) was added to MMF media for 12 h.	Sigma		Suggestion: sigma-tau (RRID:SCR_000488)(link)
The number of PC1 particles colocalized with LAMP1 was analyzed using thresholded images in Fiji.	Fiji		Suggestion: Fiji (RRID:SCR_002285)(link)
Statistical analyses GraphPad Prism software (v6.0) was used for graphs and statistics.	GraphPad Prism		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
To generate P-values for Gaussian distributed data groups of three of more, a one- or two-way ordinary ANOVA was used with Tukeys posttest or Sidaks multiple comparisons test, respectively.	ANOVA
	Tukeys posttest
	Sidaks multiple comparisons test
For pairwise comparisons of Gaussian distributed data, a Students t-test was used.	t-test
For nonparametric distributed data groups of three or more, a nonparametric ANOVA and Dunns multiple comparisons test was used.	Dunns multiple comparisons test
For pairwise comparisons of nonparametric distributed data, a Mann-Whitney test was used.	Mann-Whitney test
Oligonucleotides	
Primers used were as follows: col1a1 (forward, GCTCCTCTTAGGGGCCACT reverse, CCACGTCTCACCATTGGGG), lox (forward, GCCGACCAAGATATTCCTGGG reverse, GCAGGTCATAGTGGCTAAACT), loxl2 (forward, GGTGGAGGTGTACTATGATGG reverse, CTTGCCGTAGGAGGAGCT), tgfb1 (forward, CTCCCGTGGCTTCTAGTGC reverse, GCCTTAGTTTGGACAGGATCTG), and rpl7 to normalize gene expression (forward, ACCGCACTGAGATTCGGATG reverse, GAACCTTACGAACCTTTGGGC).	GCTCCTCTTAGGGGCCACT
	GCCGACCAAGATATTCCTGGG
	GCAGGTCATAGTGGCTAAACT
	GGTGGAGGTGTACTATGATGG
	CTCCCGTGGCTTCTAGTGC
	GCCTTAGTTTGGACAGGATCTG
	GAACCTTACGAACCTTTGGGC

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq