Document Identifier: **PMC8250065**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: Materials and methods Blood sampling Following written informed consent peripheral blood samples were obtained by vene puncturefrom peripheral venous catheters.
Inclusion and Exclusion Criteria
Inclusionexclusion criteria were: Fully conversant in the German or English language, able and willing to attend to the study requirements, written informed consent to participate in this studyhistory of four or more infections with antibiotic treatment in the year preceding the blood draw, severe liver or kidney disease, patients undergoing immunotherapy, and cancer patients.
Attrition
After the onset of symptoms all but one patient underwent RT-PCR testing and tested positive for COVID-19.
Sex as a biological variable
SARS-CoV-2-infected group: COVID-19 patients (n = 13 females, age range 23 to 70 years n = 11 male, age range 35 to 70 years) were recruited across the spectrum of clinical severity, ranging from asymptomaticmild to severeSARS.
Subject Demographics
Age: SARS-CoV-2-infected group: COVID-19 patients (n = 13 females, age range 23 to 70 years n = 11 male, age range 35 to 70 years) were recruited across the spectrum of clinical severity, ranging from asymptomaticmild to severeSARS.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication

Each peptide was deposited on the microarray in triplicates .
Number: Each peptide was deposited on the microarray in triplicates .
Cell Line Authentication
not detected.
Data Information
Availability: Each peptide was deposited on the microarray in triplicates .

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Peptide microarrays Antibody response profiling was performed with a peptide microarray (JPT Peptide Technologies GmbH) containing 5513 peptides covering the full proteome of SARS-CoV-2 and , in addition , the S , N , E , and M proteins of the following HCoVs: SARS-CoV-1 , MERS-CoV , HCoV-229E , HCoV-OC43 .	HCoV-OC43		
Microarray assay and data analysis The peptide microarrays were incubated with sera (applied dilution 1:200) in an HS 4800 microarray processing station (Tecan) for 2 h at 30degC , followed by incubation with 0.1 mgmL fluorescently labelled anti human IgG detection antibody (Jackson Immunoresearch , 109-605-098)	anti human IgG		
ELISA test To detect antibodies , we used the Euroimmun Anti-SARS-CoV-2 ELISA (IgG) , which was manufactured by Euroimmun Medizinische Labordiagnostika AG , Lubeck , Germany .	Anti-SARS-CoV-2 ELISA (IgG		
First , the diluted patient samples were incubated in the wells which lead to specific IgG antibodies binding to the antigens .	specific IgG		
In order to detect the bound antibodies , an enzyme-labelled anti-human IgG antibody detected antigen-antibody complexes and catalyzed a color reaction .	anti-human IgG		
Experimental Models: Cell Lines			
Peptide microarrays Antibody response profiling was performed with a peptide microarray (JPT Peptide Technologies GmbH) containing 5513 peptides covering the full proteome of SARS-CoV-2 and , in addition , the S , N , E , and M proteins of the following HCoVs: SARS-CoV-1 , MERS-CoV , HCoV-229E , HCoV-OC43 .	HCoV-229E		
Software and Algorithms			

http:www.r-project.org).	http:www.r-project.org		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
RT-PCR test We used the protocol described by the centers for disease control and prevention for the CDC 2019 Novel Coronavirus (2019-nCoV	RT-PCR test
Statistical comparison of two sample groups was done by the Wilcoxon Rank Sum test (R package stats, n = 24 patient samples and n = 12 control samples).	Wilcoxon Rank Sum test
The Spearmans rank correlation coefficient reflecting the relationship between the commercial ELISA and the peptide microarray (Fig. 3) was calculated as described in the figure legend using the R package ggpubr.	Spearmans rank correlation

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq