Document Identifier: **PMC8267442**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Field Sample Permit: 2.2 Collection , identification , and authentication of plant material The aerial parts of Zanthoxylum zanthoxyloides were collected from Achimota Forest , a suburb of Achimota in the Greater Accra Region of Ghana , on 4th June 2018 . Mr . Francis Otoo , the Curator at the herbarium unit , School of Biological Sciences , University of Cape Coast , Ghana , identified and authenticated the plant .
IRB: 2.6 Ethical clearance All animal experiments , procedures , and techniques used in this study were approved by the University of Cape Coast Institutional Review Board (UCCIRB) with an approval number , UCCIRBCHAS201813 .
Inclusion and Exclusion Criteria
not detected.
Attrition
Five other rats were randomly selected seven days post-diabetes induction and their blood glucose concentration tested again to confirm the establishment of type-2 diabetes .
Sex as a biological variable
2.5 Animal acquisition and husbandry Healthy 10-12 weeks old male Sprague-Dawley rats with an average weight of 182.16 +- 31.64 g) were used for the preliminary glucose kinetics effect of the ZZCE and 7-8 weeks old weighing between 130 and 160 g for the rest of the experiments .
Subject Demographics
Age: 2.5 Animal acquisition and husbandry Healthy 10-12 weeks old male Sprague-Dawley rats with an average weight of 182.16 +- 31.64 g) were used for the preliminary glucose kinetics effect of the ZZCE and 7-8 weeks old weighing between 130 and 160 g for the rest of the experiments .
Weight: 2.5 Animal acquisition and husbandry Healthy 10-12 weeks old male Sprague-Dawley rats with an average weight of 182.16 +- 31.64 g) were used for the preliminary glucose kinetics effect of the ZZCE and 7-8 weeks old weighing between 130 and 160 g for the rest of the experiments .
Randomization
Five other rats were randomly selected seven days post-diabetes induction and their blood glucose concentration tested again to confirm the establishment of type-2 diabetes .
Blinding

not detected.
Power Analysis
not detected.
Replication
not required.
Data Information
Availability: A voucher specimen (SCSBSUCCB41SH) was deposited in the herbarium for future reference.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
50 mL of diluted serum , and then 50 mL of HRP conjugated anti-Rat Insulin antibody was sequentially added to the test sample wells .	anti-Rat Insulin		
Experimental Models: Organisms/Strains			
2.5 Animal acquisition and husbandry Healthy 10-12 weeks old male Sprague-Dawley rats with an average weight of 182.16 +- 31.64 g) were used for the preliminary glucose kinetics effect of the ZZCE and 7-8 weeks old weighing between 130 and 160 g for the rest of the experiments .	Sprague-Dawley		Suggestion: RRID:RGD_70508(link)
Software and Algorithms			
The mean cross-sectional area of islets of Langerhans was estimated with ImageJ java image processing program v .	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
2.18 Statistical analysis Data were analyzed using GraphPad Prism v8.0 (	GraphPad Prism		Suggestion: GraphPad Prism (RRID:SCR_002798)(link)
(GraphPad, La Jolla, CA, USA).	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
The identity of the extract was confirmed with Mayers test and Wagners test and named Zanthoxylum zanthoxyloides alkaloidal extract (ZZAE) .	Mayers test
The data were reported as means +- SD of each group and statistical significance determined using analysis of variance (ANOVA).	analysis of variance
	ANOVA

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq