Document Identifier: **PMC8280806**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not required.
Inclusion and Exclusion Criteria
not required.
Attrition
not required.
Sex as a biological variable
not required.
Subject Demographics
Age: not required.
Weight: not required.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
Each experiment was replicated two times for Fig EV5B-C, and one time for Fig EV5D-E, using the same method with the proteins produced by different TNT reactions and the mitochondria isolated from the same group of mice.

Number: Each experiment was replicated two times for Fig EV5B-C, and one time for Fig EV5D-E, using the same method with the proteins produced by different TNT reactions and the mitochondria isolated from the same group of mice.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
After 3 min of mixing and equilibrating the sample to 37degC , the CB fluorescence intensity F(t) was measured for 3 s every 5 min for 480 min or after 480-min incubation to monitor the quenching of the released CB-dextrans by the anti-CB antibodies located outside of the liposomes .	anti-CB		
The fraction of CB-dextran release caused by the proteins equals to the extent of fluorescence quenching by the anti-CB antibody in the presence of the proteins normalized to that in the presence of Triton , i.e. , F0-F(t)F0-FT .	F0-F(t)F0-FT		
Recombinant DNA			
The resulting Bax (a2-a5) DNA was fused with a DNA encoding a His8-tag followed by a 3C protease cleavage site (LEVLFQGP) and cloned into a pET28a (	pET28a		Suggestion: RRID:Addgene_26094(link)
Plasmids encoding mutant human Bax proteins were generated from the pTYB1 vector-based plasmid encoding the wild-type protein (Suzuki etal , 2000) using QuickChange (Thermo Fisher Scientific) .	pTYB1		Suggestion: RRID:Addgene_37225(link)
Software and Algorithms			
The data were analyzed by ASTRA software , and the molecular weight of Bax (a2-a5) peak fraction was calculated using the three-detector method.	ASTRA		Suggestion: ASTRA (RRID:SCR_016255)(link)

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq