Document Identifier: **PMC8283774**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: EXPERIMENTAL MODEL AND SUBJECT DETAILS Animal Studies Animals were cared for in accordance with the NIH Guide for the Care and Use of Laboratory Animals , and all protocols were approved by the Institutional Animal Care and Use Committee (IACUC) of Tufts University School of Medicine.
Inclusion and Exclusion Criteria
not detected.
Attrition
Two of the screws served as EEG leads in the frontal cortex , 1 as a reference ground , and 1 as an animal ground.
Sex as a biological variable
All experiments save cell culture immunocytochemistry (ICC) and co-immunoprecipitation employed only male mice , aged 8-12 weeks .
Subject Demographics
Age: All experiments save cell culture immunocytochemistry (ICC) and co-immunoprecipitation employed only male mice , aged 8-12 weeks .
Randomization
not detected.
Blinding
Images were taken at a 512 x 512 resolution (17 nmpixel) with a 60x (oil , numerical aperture: 1.00) objective , with the imager blind to genotype.
Power Analysis
Baseline analysis Power analysis was performed using LabChart software v.7.3.8
Replication
not required.

Cell Line Authentication	
not detected.	
Data Information	
Identifiers: Cells were washed with PBS 10 times and mounted on glass slides (Electron Microscopy Sciences , 71864) using ProLong Gold Antifade mounting media (ThermoFisher , P36930) .	P36930

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
These PM lysates (5000 mg protein) were incubated with 150 mL Protein A Dynabeads (ThermoFisher , 10001D) cross-linked to 15 mg of N-terminal a1 antibody (Abcam , ab33299) overnight at 4*C .	a1		
Experimental Models: Cell Lines			
Graphs were plotted as mean +- SEM. Immunocytochemistry (ICC) DIV21 primary hippocampal WT and Gabra1-2 cells were washed once in PBS and fixed for 20 minutes in a solution of 4 (wv) sucrose and 4 (vv) paraformaldehyde in PBS.	Gabra1-2		
Experimental Models: Organisms/Strains			
After founder Gabra1-2 mice were received from Genoway , mice were backcrossed onto the C57Bl6J line for at least 10 generations prior to experiments .	C57Bl6J		
Western blotting 8-12-week-old male WT and Gabra1-2 mice were anaesthetized with isofluorane and their brains were rapidly removed.	Gabra1-2		
Software and Algorithms			
Images were acquired with a Nikon A1 confocal scanning laser microscope coupled with NIS Elements software v.4.20 (Nikon) .	NIS Elements		Suggestion: NIS-Elements (RRID:SCR_014329)(link)
Immunostaining density (the number of puncta per 10 mm of neuritesoma) and area were quantified using MetaMorph software (Molecular Devices) .	MetaMorph		Suggestion: MetaMorph Microscopy Automation and Image Analysis Software (RRID:SCR_002368)(link)
Co-localization analysis (calculation of the Pearsons correlation coefficient) between single-channel images was achieved with the Coloc2 macro on ImageJ v.	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
Results were analyzed on GraphPad Prism v.	GraphPad Prism		
Baseline analysis Power analysis was performed using LabChart software v.7.3.8	LabChart		
Seizure analysis The latency to the first seizure event and the latency to the onset of status epilepticus (SE)	pClamp		Suggestion: pClamp Software, RRID:SCR_011323(link)

was measured with pClamp software v.10.3 .			

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Unpaired t tests with Welchs correction were performed on GraphPad Prism v.	t tests
Co-localization analysis (calculation of the Pearsons correlation coefficient) between single-channel images was achieved with the Coloc2 macro on ImageJ v.	Pearsons correlation
Statistical analysis for mean mIPSC kinetics was performed using a Mann-Whitney test , where p < 0.05 is considered significant .	Mann-Whitney test
Tonic current was analyzed with a Students t test , where p < 0.05 is considered significant .	Students t test
All results were analyzed on GraphPad Prism v.7.10 with one-way analysis of variance (ANOVA) , with posthoc testing comparing the means between WT and Gabra1-2 hetero- and homozygotes .	analysis of variance
	ANOVA
Mortality data was plotted as a survival curve and analyzed with a log-rank (Mantel-Cox) test .	log-rank
	Mantel-Cox) test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq