Document Identifier: **PMC8284273**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: All procedures were performed in accordance with the Guide for the Care and Use of Laboratory Animals (NIH Publications No. 8023, revised 1978), and were approved by the Animal Ethics Committee of Nanfang Hospital, Southern Medical University, China (approval No. NFYY-2017-67) on June 11, 2017.
Euthanasia Agents: Twelve weeks following injury, animals were deeply anesthetized by intraperitoneal injection of sodium pentobarbital (80 mgkg).
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
Materials and Methods Animals A total of 32 adult male Sprague-Dawley rats (weighing 320 g, aged 3-4 months) were purchased from Guangdong Medical Laboratory Animal Center, Foshan, China (license No. SYXK (Yue) 2018-0002), and were randomly assigned to one of the following three groups: 1.2-mm SCI (n = 12, defined as mild), 2-mm SCI (n = 12, defined as severe), or sham (laminectomy, n = 8).
Subject Demographics
Weight: Materials and Methods Animals A total of 32 adult male Sprague-Dawley rats (weighing 320 g, aged 3-4 months) were purchased from Guangdong Medical Laboratory Animal Center, Foshan, China (license No. SYXK (Yue) 2018-0002), and were randomly assigned to one of the following three groups: 1.2-mm SCI (n = 12, defined as mild), 2-mm SCI (n = 12, defined as severe), or sham (laminectomy, n = 8).
Age: Materials and Methods Animals A total of 32 adult male Sprague-Dawley rats (weighing 320 g, aged 3-4 months) were purchased from Guangdong Medical Laboratory Animal Center, Foshan, China (license No. SYXK (Yue) 2018-0002), and were randomly assigned to one of the following three groups: 1.2-mm SCI (n = 12, defined as mild), 2-mm SCI (n = 12, defined as severe), or sham (laminectomy, n = 8).
Randomization
Materials and Methods Animals A total of 32 adult male Sprague-Dawley rats (weighing 320 g, aged 3-4 months) were purchased from Guangdong Medical Laboratory Animal Center, Foshan, China (license No. SYXK (Yue) 2018-0002), and were randomly assigned to one of the following three groups: 1.2-mm SCI (n = 12, defined as mild), 2-mm SCI (n = 12, defined as severe), or sham (laminectomy, n = 8).

Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Organisms/Strains			
Materials and Methods Animals A total of 32 adult male Sprague-Dawley rats (weighing 320 g, aged 3-4 months) were purchased from Guangdong Medical Laboratory Animal Center, Foshan, China (license No. SYXK (Yue) 2018-0002), and were randomly assigned to one of the following three groups: 1.2-mm SCI (n = 12, defined as mild), 2-mm SCI (n = 12, defined as severe), or sham (laminectomy, n = 8).	Sprague-Dawley		Suggestion: RRID:RGD_70508(link)
Software and Algorithms			
The damaged area of the spinal cord was measured using ImageJ software (National Institutes of Health, Bethesda, MD, USA).	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
Statistical analysis All statistical analyses were performed using SPSS 20.0 (IBM, Armonk, NY, USA).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Between-group differences in SEP and MEP parameters were assessed using a multiple factor repeated-measures analysis of variance followed by Tukeys honestly significant test.	Tukeys honestly significant test
Following the one-way analysis of variance, Bonferroni post hoc tests between groups were used to analyze the tissue lesion area and spared white and gray matter areas.	analysis of variance
	Bonferroni
An independent linear correlation analysis (Pearsons correlation coefficient) was performed between the individual behavioral outcomes and electrophysiological parameters.	Pearsons correlation

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq