Document Identifier: **PMC8286250**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not detected.
Inclusion and Exclusion Criteria
not required.
Attrition
not required.
Sex as a biological variable
Materials and Methods Animals C57BL6 mice (pathogen-free, male, age 5-6 weeks) were purchased from the SLRC Laboratory (Shanghai, China), and housed in controlled conditions of temperature (21-25 degC) and humidity (40-60), with ad libitum access to water and food (free from OVA).
Subject Demographics
Age: Materials and Methods Animals C57BL6 mice (pathogen-free, male, age 5-6 weeks) were purchased from the SLRC Laboratory (Shanghai, China), and housed in controlled conditions of temperature (21-25 degC) and humidity (40-60), with ad libitum access to water and food (free from OVA).
Randomization
Allergic Airway Inflammation Induction and Ozone Exposure C57BL6 mice were randomly divided into three groups with eight mice in each group (see schematic diagram of mouse experiment in Figure S2 of Supplementary information).
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Immunohistochemical Analyses Airway smooth muscle (ASM) mass and airway mucus secretion was observed using immunohistochemical staining, based on the presence of alpha-smooth muscle actin (a-SMA) and Muc5ac protein in lung tissue sections with the mouse a-SMA antibody (Proteintech, Wuhan, China) and mouse Muc5ac antibody (GTX11335, GeneTex, USA), as previously described.	Airway smooth muscle (ASM		
	alpha-smooth muscle actin (a-SMA)		
	a-SMA		
	Muc5ac		
Experimental Models: Organisms/Strains			
Allergic Airway Inflammation Induction and Ozone Exposure C57BL6 mice were randomly divided into three groups with eight mice in each group (see schematic diagram of mouse experiment in Figure S2 of Supplementary information).	C57BL6		Suggestion: RRID:IMSR_JAX:000664(link)
Software and Algorithms			
16 The area of a-SMA-positive and Muc5ac-positive expression around the airway epithelial cells in the respective images were independently analyzed in a double-blind fashion by two investigators, using the Image J software.	Image J		Suggestion: ImageJ, RRID:SCR_003070(link)
Statistical Analysis Statistical analysis was conducted and graphs were constructed using the GraphPad PRISM, version 5.0 software (GraphPad, San Diego, CA).	GraphPad PRISM		
	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
The Spearman correlation coefficient matrix was constructed and Spearman rank correlation tests were performed using the R version 3.6.1 (Innovative Solutions, St. Louis, MO, USA).	Spearman correlation
	Spearman rank correlation
The Kruskal-Wallis ANOVA with Bonferronis post hoc test (for equal variance) or Dunnetts T3 post hoc test (for unequal variance) were used to evaluate the differences in variance between multiple groups.	Kruskal-Wallis ANOVA
	Bonferronis post hoc test
	Dunnetts T3 post hoc test
A posttest Mann-Whitney analysis was conducted to evaluate the differences in variance between two groups.	Mann-Whitney
The correlation between different variables was determined using Spearman analysis $r > 0.4$ or < -0.4 was considered indicative of strong correlations, and r values between -0.4 and 0.4 were considered indicative of weak correlations if $P < 0.05$. $P < 0.05$ was considered statistically significant.	Spearman
Oligonucleotides	
The following primer sequences were used in the PCR: b-actin: Forward-5 CCT CTA TGC CAA CAC AGT 3, Reverse-5 AGC CAC CAA TCC ACA CAG 3 Muc5ac: Forward-5 TCA CAT TTG ATG GCA CCT ACT 3, Reverse-5 CAC ACG GAA GTA TCC AAA CAC 3 Histological and Morphometric Analysis After the lungs of mice were resected, the left lungs were inflated by injection of 4 paraformaldehyde to provide 20 cm of water pressure.	CCT CTA TGC CAA CAC AGT 3
	AGC CAC CAA TCC ACA CAG 3
	TCA CAT TTG ATG GCA CCT ACT 3
	CAC ACG GAA GTA TCC AAA CAC 3

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq