Document Identifier: **PMC8292958**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: This study was approved by the Ethics Committee of aforementioned hospital.
Consent: All patients signed the informed consent.
Inclusion and Exclusion Criteria
All patients were newly diagnosed cases and patients with other clinical disorders were excluded.
Attrition
These patients were grouped into 27 cases at AJCC stage I or II, and 37 cases at AJCC stage III or IV.
Sex as a biological variable
Materials and Methods Tissue Acquisition HCC and paired adjacent (within 3 cm around tumors) non-tumor tissues were collected from a total of 64 HCC patients (40 males and 24 females) who were admitted at Xuzhou Cancer Hospital between May 2018 and May 2020.
Subject Demographics
Age: The age range of patients was 44–68 years old, with a median age of 55.8 years old.
Randomization
Migration cells were observed and counted in 3 randomly selected fields per well under a light microscope as previously described.
Blinding
not detected.
Power Analysis
not detected.
Replication
All reactions were performed in 3 technical replicates .
Type: All reactions were performed in 3 technical replicates .

Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Cell Lines			
HCC Cells and Transfections Two human HCC cell lines SNU-423 and SNU-387 (ATCC, USA) were used as the cell models of HCC.	SNU-423		Suggestion: (RRID:CVCL_0366)(link)
	SNU-387		Suggestion: (RRID:CVCL_0250)(link)
HCC cells were seeded into a 6-well plate with 1.0×106 cells/well , followed by transfection with 1 ug vector and 40 nM synthetic miRNA oligonucleotides using Lipofectamine 2000 (Invitrogen) .	HCC		
Recombinant DNA			
To overexpress circRNA SCARB1, pcDNA3.1(	pcDNA3.1		Suggestion: RRID:Addgene_100544(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Data comparisons were performed by paired t-test.	t-test
Data comparisons were performed by ANOVA Tukey's test.	Tukey's test
Correlations were analyzed by Pearson's correlation coefficient.	Pearson's correlation
Oligonucleotides	
The sequences of primers used in qRT-PCR assay were shown as follows: CircRNA SCARB1 (Forward: 5#-TCATCCTTCTAAGGTAGCAGC AG-3#; Reverse: 5#-CTTCGACGCTGTAGCCTTGT-3#); mature miR-497 (Forward: 5#-CCTTCAGCAGCACACTGTGG- 3#;	5#-TCATCCTTCTAAGGTAGCAGC AG-3#
	5#-CTTCGACGCTGTAGCCTTGT- 3#)
	5#-CCTTCAGCAGCACACTGTGG- 3#
miR-497 precursor (Forward: 5#-ACCAGCAGCACACTGTGGTTT GT-3#; Reverse: 5#-ATCCAGTGCAGGGTCCGAGG- 3#) .	5#-ACCAGCAGCACACTGTGGTTT GT-3#
	5#-ATCCAGTGCAGGGTCCGAGG- 3#

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq