Document Identifier: **PMC8299063**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not detected.
Inclusion and Exclusion Criteria
not required.
Attrition
not required.
Sex as a biological variable
not detected.
Subject Demographics
Age: Each pGWB605 construct or the combinations of YFPN and YFPC fusion constructs were co-expressed in 6- to 8-week-old Nicotiana benthamiana leaves after leaf infiltration with GV3101 strains containing the tested construct combinations plus an anti-silencing Agrobacterium strain expressing P19 .
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
Three technical replicates were performed for each strain in each growth assay .
Type: Three technical replicates were performed for each strain in each growth assay .
Number: Three technical replicates were performed for each strain in each growth assay .

Cell Line Authentication
Authentication: Homozygous lines were identified by genotyping .

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Cell Lines			
Y2H assays Plasmid vectors pGAD424 and pGBT9 (Clontech) containing the cDNAs of BRM , SWI3A , SWI3B , BSH , and SWP73B were described previously (Buszewicz et al. , 2016) .	SWI3A		
	SWI3B		
	SWP73B		
The full-length cDNAs of BRD1 , BRD2 , BRD13 , SWI3C , SWI3D , SWP73A , ARP4 , and ARP7 and a cDNA encoding the N-terminal region of SYD (1-1500 bp of the coding sequence) were cloned into pGAD424 and pGBT9 by ligation with T4 DNA ligase (Thermo Scientific) or using the SLIC protocol (Li and Elledge , 2007) .	SWP73A		
Experimental Models: Organisms/Strains			
The brm-1BRM-GFP and brdx3 brm-1BRM-GFP lines were constructed by crossing the pBRM:BRM-GFP line with the respective mutants .	brdx3 brm-1BRM-GFP		
brm-1 brdx3 mutants were identified from segregating progeny of the brdx3 brm-1BRM-GFP +- line.	brdx3 brm-1BRM-GFP +-		
Recombinant DNA			
The brm-1BRM-GFP and brdx3 brm-1BRM-GFP lines were constructed by crossing the pBRM:BRM-GFP line with the respective mutants .	pBRM:BRM-GFP		
Y2H assays Plasmid vectors pGAD424 and pGBT9 (Clontech) containing the cDNAs of BRM , SWI3A , SWI3B , BSH , and SWP73B were described previously (Buszewicz et al. , 2016) .	pGBT9		Suggestion: RRID:Addgene_198526(link)
The full-length cDNAs of BRD1 , BRD2 , BRD13 , SWI3C , SWI3D , SWP73A , ARP4 , and ARP7 and a cDNA encoding the N-terminal region of SYD (1-1500 bp of the coding sequence) were cloned into pGAD424 and pGBT9 by ligation with T4 DNA ligase (Thermo Scientific) or using the SLIC protocol (Li and Elledge , 2007) .	pGAD424		

To generate these constructs , the full-length coding sequence of each gene was PCR-amplified using primers encoding an SV40 NLS added to the N terminus to ensure nuclear localization in yeast cells (Sun et al. , 2011) and cloned into the BamHISalI sites of the plasmid p426 carrying the URA3 nutritional marker to permit selection on uracil-deficient medium (Mumberg et al. , 1995)	p426		
BiFC and localization analyses Full-length BRD1 , BRD2 , BRD13 , and SWI3C cDNAs were cloned into the pDONR201 vector and verified by sequencing .	pDONR201		
Corresponding entry clones were used in LR recombination reactions to transfer the DNA fragments into the gateway-compatible expression vectors pYFN43 and pYFC43 for BiFC assays (Belda-Palazon et al. , 2012) or the pGWB series vector pGWB605 (Nakamura et al. , 2010) for localization analyses .	pYFN43		
	pGWB		
	pGWB605		
Software and Algorithms			
Data analysis was performed using Bioconductor packages in the R environment .	Bioconductor		Suggestion: Bioconductor (RRID:SCR_006442)(link)
Differential gene expression analysis was performed with the limma package (Ritchie et al. , 2015) .	limma		Suggestion: LIMMA (RRID:SCR_010943)(link)
The AgriGO online tool was used for GO analysis (http:systemsbiology.cau.edu.cnagriGOv2) (Tian et al. , 2017)	AgriGO		
The relative BRM signal was quantified with ImageJ software by comparison with the protein level on the Coomassie-stained gel (loading control) .	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
Phylogenetic analysis Full-length amino acid sequences of yeast and human SWISNF subunits containing bromodomains, Arabidopsis BRD1, BRD2, and BRD13, and other proteins representing previously described classes of Arabidopsis bromodomain-containing proteins (Rao et al., 2014) were used for multiple sequence alignment with ClustalW (supplemental Table 4) (Larkin et al., 2007).	ClustalW		
A neighbor-joining tree was constructed using MEGA7 software	MEGA7		

(Kumar et al., 2016), and bootstrap probability values were obtained from 1000 iterations.			
Bromodomains were detected using the ScanProsite tool from the Prosite database.	Prosite		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Data were fitted to a linear model , and moderated t-statistics were computed by empirical Bayes moderation .	t-statistics

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq