Document Identifier: **PMC8320599**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: Ethical approval All study protocols were reviewed and approved by the Institutional Animal Care and Use Committee of Texas A&M University. 2.2.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
Study 1 (global Pfkfb3 disruption on NAFLD phenotype): male Pfkfb3 +- mice and WT littermates, at 5-6 weeks of age, were fed an HFD (60 fat calories, 20 carbohydrate calories, and 20 protein calories Research Diets, Inc.) or low-fat diet (LFD,10 fat calories, 70 carbohydrate calories, and 20 protein calories) for 12 weeks as previously described.
Subject Demographics
Age: Study 1 (global Pfkfb3 disruption on NAFLD phenotype): male Pfkfb3 +- mice and WT littermates, at 5-6 weeks of age, were fed an HFD (60 fat calories, 20 carbohydrate calories, and 20 protein calories Research Diets, Inc.) or low-fat diet (LFD,10 fat calories, 70 carbohydrate calories, and 20 protein calories) for 12 weeks as previously described.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Data Information	
Identifiers: 22,25,26 Study 2 (hematopoietic cell-specific Pfkfb3 disruption on NAFLD phenotype): to generate chimeric mice in which Pfkfb3 was disrupted only in hematopoietic cells, male WT mice, at 5-6 weeks of age, were subjected to bone marrow transplantation (BMT) as described below.	22,25,26

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Histological and immunohistochemical analyses Paraffin-embedded mouse liver blocks were cut into sections of 5 mm thickness and stained with hematoxylin and eosin (H&E) andor stained for F480 expression with rabbit anti-F480 antibodies (1:100) (AbD Serotec, Raleigh, NC, USA). 6,23 The fraction of F480-expressing cells for each sample is calculated as the sum of the number of nuclei of F480-expressing cells divided by the total number of nuclei in sections of a sample.	anti-F480		
Experimental Models: Organisms/Strains			
Animal experiments Wild-type (WT) C57BL6J mice were obtained from The Jackson Laboratory (Bar Harbor, ME, USA).	C57BL6J		Suggestion: RRID:IMSR_JAX:000664(link)
Software and Algorithms			
The maximum intensity of each band was quantified using Image J software.	Image J		Suggestion: ImageJ, RRID:SCR_003070(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical significance was determined using unpaired, two-tailed ANOVA or Students t tests.	ANOVA
	Students t tests

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq