Document Identifier: **PMC8320644**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: The project was filed under number 1089 with the Ethics Committee on Animal Experimentation of the UNESP Institute of Biosciences, in Botucatu.
Euthanasia Agents: On GD 20, females were weighed and euthanized by CO2.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
2 Material and methods 2.1 Animals Adult male and female Wistar rats (80 days old) were obtained from the Central Biotherium, Sao Paulo State University (UNESP), campus at BotucatuSP, Brazil, and maintained under controlled conditions (12 h of light12 h of darkness average temperature of 23 degC) in the Small Mammal Biotherium at the UNESP Department of Structural and Functional Biology (Morphology), Institute of Biosciences in Botucatu, with food and water ad libitum.
Subject Demographics
Age: 2 Material and methods 2.1 Animals Adult male and female Wistar rats (80 days old) were obtained from the Central Biotherium, Sao Paulo State University (UNESP), campus at BotucatuSP, Brazil, and maintained under controlled conditions (12 h of light12 h of darkness average temperature of 23 degC) in the Small Mammal Biotherium at the UNESP Department of Structural and Functional Biology (Morphology), Institute of Biosciences in Botucatu, with food and water ad libitum.
Randomization
2.2 Experimental design At weaning, on PND 21, female pups were randomly distributed among three experimental groups: control group, which received treatment with saline and rosuvastatin groups, which received the statin at doses of 3 or 10 mgKg.bw-1 (body weight) diluted in saline.
Blinding
not detected.
Power Analysis

not detected.
Replication
not required.
Data Information
Availability: The fluid was deposited on a slide and analyzed under light microscopy.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Serum levels of FSH, LH, prolactin were measured using a double-antibody radioimmunoassay with specific kits provided by the National Hormone and Peptide Program (Harbor-University of California at Los Angeles).	LH		
In the next step, the sections were incubated for 30 min with Bovine Serum Albumin (BSA 3) diluted in PBS, and then washed with PBS to be incubated overnight with the primary anti-PCNA antibody (PCNA PC10: sc-56, Monoclonal, Santa Cruz Biotechnology, CA, USA - 1:100).	anti-PCNA		
After the incubation period, the sections were washed again with PBS and incubated for 1 h with the secondary antibody (Goat Anti-Mouse peroxidase-labeled IgG, Catalog No. 474-1806	Anti-Mouse peroxidase-labeled IgG		
Experimental Models: Organisms/Strains			
2 Material and methods 2.1 Animals Adult male and female Wistar rats (80 days old) were obtained from the Central Biotherium, Sao Paulo State University (UNESP), campus at BotucatuSP, Brazil, and maintained under controlled conditions (12 h of light12 h of darkness average temperature of 23 degC) in the Small Mammal Biotherium at the UNESP Department of Structural and Functional Biology (Morphology), Institute of Biosciences in Botucatu, with food and water ad libitum.	Wistar		Suggestion: RRID:RGD_13508588(link)
Software and Algorithms			
Histological analyses were conducted under light microscopy, with the software packages Leica QWin 3 and ImageJ 1.48.	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
Statistical analyses were performed using the software GraphPad Prism (Version 6.0).	GraphPad Prism		Suggestion: GraphPad Prism (RRID:SCR_002798)(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Shapiro-Wilks normality test was applied in the intragroup values, then results were compared among groups by one-way analysis of variance (ANOVA) followed by Tukeys test, for parametric variables, and by Kruskal-Wallis followed by Dunns test, for nonparametric variables.	Shapiro-Wilks normality test
	analysis of variance
	ANOVA
	Tukeys test
	Kruskal-Wallis
	Dunns test
Then, the proportions were analyzed by the Fisher exact test.	Fisher exact test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq