Document Identifier: **PMC8328822**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not detected.
Inclusion and Exclusion Criteria
not required.
Attrition
not required.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Mass spectrometry Rat brain lysate was immunoprecipitated with anti-βPIX antibody immobilized onto CNBr-activated sepharose 4B .	anti-βPIX		
GST-βPIX-immo-bilized beads were incubated with His-tagged FL-KIF5A, motor (aa 1-330), stalk (aa 331-799), or tail (aa 800-1027) in binding buffer (20 mM HEPES, pH 7.4, 0.15 M NaCl, 1 mM DTT, 0.2% Triton X-100, 5 mM MgSO4, and protease inhibitor) for 1 h, and then subjected to immunoblotting with antibodies to His and GST.	GST		
Experimental Models: Cell Lines			
MATERIALS AND METHODS Cell culture PC12 cells were grown in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS) and 100 U/ml penicillin/streptomycin (Invitrogen) and were kept at 37°C in a humidified 5% CO2 incubator.	PC12		Suggestion: RRID:CVCL_0481(link)
Experimental Models: Organisms/Strains			
Hippocampus from the brain of newborn (P1) Sprague-Dawley (SD) rat were dissected and incubated with papain in dissociation medium (pH 7.4) (82 mM Na2SO4, 30 mM K2SO4, 5.8 mM MgCl2, 0.252 mM CaCl2, 1 mM HEPES, 20 mM glucose, 0.5% phenol red) containing L-cystein for 20 min at 37°C and dissociated by pipetting in DMEM containing 10	Sprague-Dawley		Suggestion: RRID:RGD_70508(link)
Recombinant DNA			
In vitro binding assay pGEX4T-1-βPIX and pET-FL-KIF5 constructs were expressed in E. coli DH5α or BL21.	pGEX4T-1-βPIX		
	pET-FL-KIF5		
Software and Algorithms			
Statistical significance was assessed using an unpaired t-test (t-test) using GraphPad Prizm software (version 10) for Windows.	GraphPad Prizm		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical significance was assessed using an unpaired t-test (t-test) using GraphPad Prizm software (version 10) for Windows.	t-test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq