Document Identifier: **PMC8346374**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Euthanasia Agents: 4 Dams were fasted for 4 h and euthanized individually using CO2 asphyxiation in the afternoon (1500 h) in a room separate from where other animals were housed.
IACUC: IACUC Approval: All protocols were approved by the University of Minnesota Institutional Animal Care and Use Committee.
Inclusion and Exclusion Criteria
Litters containing fewer than five or greater than eight fetuses were excluded from fetal studies to control for variation in fetal nutrient exposure , resulting in a final cohort of 9 CON litters and 8 HFHS litters studied.
Attrition
One Sry + fetus and one Sry- fetus per litter were studied (CON: n = eight male , n = nine female HFHS: n = eight male , n = seven female) .
Sex as a biological variable
2 MATERIALS AND METHODS 2.1 Animals and diets Animals were group housed (two pregnant dams per cage) under controlled conditions (25degC, 12-h lightdark cycle, lights on 0600 h).
Subject Demographics
Age: Female, 6-week-old, C57Bl6J mice (The Jackson Laboratory) were randomly allocated to be fed ad libitum either a control diet (D12489B Research Diets, 10.6 kcal fat, 16.8 kcal protein, 72.6 kcal carbohydrate, 240 gkg sucrose) (CON group) or a HFHS diet (HFHS group).
Randomization
Female, 6-week-old, C57Bl6J mice (The Jackson Laboratory) were randomly allocated to be fed ad libitum either a control diet (D12489B Research Diets, 10.6 kcal fat, 16.8 kcal protein, 72.6 kcal carbohydrate, 240 gkg sucrose) (CON group) or a HFHS diet (HFHS group).
Blinding
not detected.
Power Analysis
not detected.

Replication
Number: Fetal sex was determined by RT-qPCR assessment of Sry gene amplification in three randomly chosen fetuses per litter (N = 51) .

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Organisms/Strains			
17 Insulin sensitivity was measured by surrogate measures of insulin resistance: homeostasis model of insulin resistance (HOMA-IR = fasting insulinngml x fasting glucosemgdL22.5) and quantitative insulin check index of insulin sensitivity (QUICKI = 1Log(insulin) + Log(glucose)) which have been previously validated in C57Bl6J mice.	C57Bl6J		
Software and Algorithms			
, Actb, Pgk1, Hprt) were evaluated in whole brain and placenta homogenates by geNorm, 20 NormFinder, 21 the comparative delta-Ct method, 22 and BestKeeper 23 using a web-based program (https:www.heartcure.com.aureffinder) 24 (Table S2).	geNorm		Suggestion: geNORM (RRID:SCR_006763)(link)
	NormFinder		Suggestion: NormFinder (RRID:SCR_003387)(link)
	BestKeeper		Suggestion: BestKeeper (RRID:SCR_003380)(link)
Statistical analysis and graphics were performed using GraphPad Prism version 7 (GraphPad Software, Inc).	GraphPad Prism		
	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
To determine sample size, the resource equation was used (Formula: E = total number of animals-total number of groups) where the degree of freedom (E) must be greater than 10. 25 , 26 , 27 Data from primary outcome groups passed the Shapiro-Wilk normality test therefore, results were analyzed using unpaired t-test to measure group differences and two-way ANOVA followed by post-hoc Tukeys HSD to measure diet and sex effects between groups.	Shapiro-Wilk
	t-test
	ANOVA
Pearson correlation coefficient was used as a measure of linear correlation.	Pearson correlation

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq