Document Identifier: **PMC8350004**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Ethical approval was obtained by the NHS Health Research Authority Blackcountry Research Ethics Committee (REF Reference 17/WM/0125) .
Consent: All participants provided written informed consent at baseline assessment .
Inclusion and Exclusion Criteria
Eligible participants were adults with a DSM/ADI-R or equivalent confirmed diagnosis of an Autism Spectrum Condition, aged 18–65 years, with normal or corrected-to-normal vision and hearing, and fluent understanding of English.
Attrition
No other adjustments for multiple testing were recommended by the trial statistician.
Sex as a biological variable
not detected.
Subject Demographics
Age: Eligible participants were adults with a DSM/ADI-R or equivalent confirmed diagnosis of an Autism Spectrum Condition, aged 18–65 years, with normal or corrected-to-normal vision and hearing, and fluent understanding of English.
Randomization
2 Methods 2.1 Study design and participants In this parallel-group, superiority, randomized controlled trial, we recruited 121 autistic adults from current and former patients of the Sussex Partnership NHS Foundation Trust Neurodevelopmental Service, and through community and third-sector organizations.
Blinding
Researchers collecting outcome data and the trial statistician were blind to treatment allocation .
Power Analysis
2.5 Sample size The sample size calculation was based on the use of an independent t-test to detect a minimal clinically meaningful difference (MCID) in the primary outcome STAI-T of 7·65 or more at the primary time point (T2) .

Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
However, a small number of secondary outcomes were essentially measuring similar constructs, i.e., STAI-S and GAD-7 (both measuring anxiety), and BPQ and MAIA (both measuring sensibility to bodily signals), and so Bonferroni corrections for multiple comparisons were applied.	MAIA		Suggestion: MAIA (RRID:SCR_007153)(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
2.5 Sample size The sample size calculation was based on the use of an independent t-test to detect a minimal clinically meaningful difference (MCID) in the primary outcome STAI-T of 7·65 or more at the primary time point (T2) .	t-test
Secondary outcomes, for which only T0 and T1 data were available, were evaluated using Analysis of Covariance (ANCOVA) at T1, with corresponding baseline scores as covariates, and treatment group (ADIE, prosody) as a fixed factor.	Analysis of Covariance
	ANCOVA
However, a small number of secondary outcomes were essentially measuring similar constructs, i.e., STAI-S and GAD-7 (both measuring anxiety), and BPQ and MAIA (both measuring sensibility to bodily signals), and so Bonferroni corrections for multiple comparisons were applied.	Bonferroni corrections

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq