Document Identifier: **PMC8350424**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: All procedures used in this study have conformed to the rules and principles of the 86609EEC Directive and have been approved by the Local Bioethical Committee at the Institute of Pharmacology , Polish Academy of Sciences , Krakow , Poland.
Euthanasia Agents: Based on the result of the sucrose consumption test carried out following the first 2 weeks of stress , two groups of animals were identified: those showing a significative reduction of sucrose intake in comparison to the control group , ie. developing an anhedonic-like behavior , (CMS vulnerable) and those that did not develop anhedonia (CMS resilient) . 24 h after the final stressor animals were decapitated (between 9:10 and 12:25 a . m .) , the trunk blood was collected in tubes with ethylenediaminetetraacetic acid and the ventral hippocampus (vHip) was dissected from the whole brain according to the plates 34-43 of the atlas of Paxinos and Watson (2004) and stored at -80 degC .
Field Sample Permit: 2.5 Measurement of corticosterone plasma levels After collection , blood specimens were centrifuged for 20 min at 3000 g at 4 degC for the separation of plasma .
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
2.3 Animals Adult male (post natal day 70) Wistar rats (Charles River , Germany) were adapted , housed in groups of 10 , to the animal facilities one month before the start of the experiment .
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
Animals were randomly divided into two groups: one group of animals was unstressed (No stress group) and the other one was exposed to the CMS procedure for a period of 2 consecutive weeks .
Blinding

not detected.
Power Analysis
not detected.
Replication
Samples were run in 384 well formats in triplicate as multiplexed reactions with the normalizing internal control 18s .
Number: Samples were run in 384 well formats in triplicate as multiplexed reactions with the normalizing internal control 18s .
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Italia . Anti-BDNF (SAB 1405514) and anti b-tubulin (T0198) were purchased from Sigma Aldrich	anti b-tubulin		
Host-specific peroxidase-conjugated IgG secondary antibodies were purchased from Thermo Scientific Inc .	Host-specific peroxidase-conjugated IgG secondary antibodies		
	Host-specific peroxidase-conjugated IgG		
The proteins were visualized using primary antibodies diluted in 5 wv BSA , 1X TBS , 0.1 Tween-20 for RACK1 (1:1000) , SRp30c (1:500) ,	RACK1		
	SRp30c		
Experimental Models: Cell Lines			
2.2 Cell cultures and treatments Human neuroblastoma SH-SY5Y cells from the European Collection of Cell Cultures (ECACC No. 94030304) were cultured in a medium with equal amounts of Eagles minimum essential medium and Nutrient Mixture Hams F-12 , supplemented with 10 heat-inactivated fetal bovine serum (FBS) , 2 mM glutamine , 0.1 mgml streptomycin , 100 IUml penicillin and 1X MEM non-essential amino acid solution (M7145 , Merck) at 37 degC in 5 CO2-containing , and 95 air atmosphere.	SH-SY5Y		Suggestion: RRID:CVCL_0019(link)
Experimental Models: Organisms/Strains			
2.3 Animals Adult male (post natal day 70) Wistar rats (Charles River , Germany) were adapted , housed in groups of 10 , to the animal facilities one month before the start of the experiment .	Wistar		Suggestion: RRID:RGD_13508588(link)
Software and Algorithms			
2 Material and methods 2.1 Chemicals, culture media and supplements Cortisol (PubChem CID:5754) and mifepristone (PubChem CID:55245) were obtained from Sigma Aldrich (St. Louis, MO, USA).	PubChem		Suggestion: PubChem (RRID:SCR_004284)(link)

Cell culture media and all supplements were obtained from Sigma Aldrich.	Sigma Aldrich		Suggestion: Sigma-Aldrich (RRID:SCR_008988)(link)
Total protein content was measured according to the Bradford Protein Assay procedure (Bio-Rad Laboratories) , using bovine serum albumin (BSA) as a calibration standard.	Bio-Rad Laboratories		
After Western blot acquisition, bands optical analysis was performed with the ImageJ program (W. Rasband, Research Service Branch, National Institute of Mental Health, National Institutes of Health, Bethesda, MD and Laboratory for Optical and Computational Instrumentation, University of Wisconsin).	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
Statistical analyses were conducted with GraphPad Prism version 7 (GraphPad Software, San Diego, CA, USA) and outliers were determined with GraphPad outlier calculator (alpha = 0.05).	GraphPad Prism		
	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical differences were determined by analysis of variance (one-way ANOVA) followed, when significant, by the Tukey multiple comparison post hoc test.	analysis of variance
	ANOVA
	Tukey multiple comparison post hoc test
Oligonucleotides	
Table 1 a) Gene Forward primer Reverse primer Probe Gra GCGACAGAAGCAGTTGAGTCATC CCATGCCTCCACGTAACTGTTAG TGAAGTGATAGCACAGCAGACAGT GA Grb GCGCTTGAGGCTAAGATAGCT CCCATGTTTCTGCCTCTTTCTTTG AGTCTGCCTTCAGAATGCCTGTCA Rack1 CTCTTTGGCTTGGTCTGCTG ATACACGCACCAAGTTGTCG	GCGACAGAAGCAGTTGAGTCATC CCATGCCTCCACGTAACTGTTAG TGAAGTGATAGCACAGCAGACAGT GA Grb GCGCTTGAGGCTAAGATAGCT CCCATGTTTCTGCCTCTTTCTTTG AGTCTGCCTTCAGAATGCCTGTCA Rack1 CTCTTTGGCTTGGTCTGCTG ATACACGCACCAAGTTGTCG
ATGGCCAGACTCTGTTTGCT Total Bdnf AAGTCTGCATTACATTCCTCGA GTTTTCTGAAAGAGGGACAGTTTA T TGTGGTTTGTTGCCGTTGCCAAG 18s GTAACCCGTTGAACCCCATT CCATCCAATCGGTAGTAGCG TGCAATTATTCCCCATGAACGAGG b) Gene Accession number Assay ID Bdnf long 3UTR EF125675	AAGTCTGCATTACATTCCTCGA GTTTTCTGAAAGAGGGACAGTTTA T TGTGGTTTGTTGCCGTTGCCAAG 18s GTAACCCGTTGAACCCCATT CCATCCAATCGGTAGTAGCG TGCAATTATTCCCCATGAACGAGG b

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq