Document Identifier: **PMC8352006**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: Informed consent was obtained, and participants and researchers were matched by gender.
IRB: Ethics approval was received from the Institutional Review Boards at Istanbul Sehir University and the London School of Hygiene and Tropical Medicine.
Inclusion and Exclusion Criteria
Individuals who were under the influence of alcohol or drugs, who faced severe mental impairments, or who were non-native Arabic speakers were excluded.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: 2 Methods 2.1 Study design A cross-sectional survey was conducted in February-May 2018 among Syrian refugees aged 18 years old and older living in Sultanbeyli, Istanbul, Turkey.
Randomization
Participants were randomly sampled using Sultanbeyli Municipalitys registration system, which maintains a record of all Syrian refugees living in Sultanbeyli granted the official Temporary Protection status from the Turkish government (Republic of Turkey: Ministry of Interior, 2020).
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Cell Line Authentication
Authentication: The PTGI has been previously validated in conflict-affected populations (Powell et al., 2003 Salo et al., 2005), and the Arabic translation has been validated in multiple Palestinian samples (Kira et al., 2012 Shamia et al., 2015 Veronese and Pepe, 2019).

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Cell Lines			
The following mental health-related measures were also screened: depression and anxiety symptoms using the Hopkins Symptoms Checklist (HSCL-25) (Mollica et al., 2004), where scores were summed and compared to a cut-off score of >=2.1 (depression) and >=2.0 (anxiety) (Fares et al., 2019) PTSD symptoms measured using the PTSD Checklist for DSM-5 (PCL-5), where scores were summed and compared to a cut-off of >=33 (Bovin et al., 2016) and somatic distress symptoms using the Patient Health Questionnaire-15 (PHQ-15), where a range of scores from 0 to 30 measured four severity distress levels of minimal (0-4), mild (5-9), moderate (10-14), or severe (>=15) (Kroenke et al., 2002).	DSM-5		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Initial T-tests and ANOVA tests were run to detect any differences in mean PTG scores between demographic and socioeconomic groups.	Initial T-tests
	ANOVA

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq