Document Identifier: **PMC8353912**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: 2.6 Animals and tumor models All in vivo procedures were conducted in compliance with the Guide for the Care and Use of Laboratory Animal Resources (1996), US National Research Council, and approved by the local Animal Care and Use Committee.
Euthanasia Agents: All in vivo procedures were performed under anesthesia using inhaled 2-5 isoflurane andor intraperitoneal injection of 0.75 mg of sodium pentobarbital (Ovation Pharmaceuticals, Deerfield, IL, USA).
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
Six- to eight-wk-old female C57BL6 mice were purchased from the Jackson Laboratory (Bar Harbor, ME, USA).
Subject Demographics
Age: Six- to eight-wk-old female C57BL6 mice were purchased from the Jackson Laboratory (Bar Harbor, ME, USA).
Randomization
RFI was calculated from mean fluorescence intensity (MFI) as (MFI of measured time point)(MFI of pre-injection status). 2.9 In vivo NIR-PIT Tumor bearing mice were randomized into 3 experimental groups: (1) no-treatment (control group), (2) intravenous (IV) injection of anti-CD44-IR700 (100 mg) without NIR light exposure (APC-IV group), and (3) IV injection of anti-CD44-IR700 (100 mg) followed by NIR light exposure (NIR-PIT group).
Blinding
not detected.
Power Analysis
not detected.

Replication
not required.
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Anti-mouse-CD44 antibody (IM7) was purchased from Bio X Cell (West Lebanon, NH, USA).	Anti-mouse-CD44		
To confirm the specific binding of the anti-CD44-IR700, a 10-fold excess of unconjugated anti-CD44 antibody was added to some samples 30 min prior to the administration of the APC.	anti-CD44-IR700		
	anti-CD44		
The cells were stained with antibodies anti-CD31(clone 390 BioLegend, San Diego, CA, USA), anti-CD44 (clone IM7 Thermo Fisher Scientific), anti-CD45 (clone 30-F11 Thermo Fisher Scientific), and anti-CD140a (clone 135 905 BioLegend).	anti-CD31		
	anti-CD45		
	anti-CD140a		
Experimental Models: Cell Lines			
2.3 Cell culture The MOC2 cell line (murine oral cancer) was purchased from Kerafast (Boston, MA, USA) and luciferase was transduced with RediFect Red-Fluc lentivirus (PerkinElmer, Waltham, MA, USA).	MOC2		Suggestion: (RRID:CVCL_ZD33)(link)
Experimental Models: Organisms/Strains			
Six- to eight-wk-old female C57BL6 mice were purchased from the Jackson Laboratory (Bar Harbor, ME, USA).	C57BL6		Suggestion: RRID:IMSR_JAX:000664(link)
Software and Algorithms			
The cells were cultured in a mixture of Iscoves Modified Dulbeccos Medium (IMDM) and Hams Nutrient Mixture F12 Medium (at a ratio of 2:1, GE Health Healthcare Life Sciences) supplemented with 5 fetal bovine serum (Thermo Fisher Scientific, Rockford, IL, USA), 1 penicillinstreptomycin (Thermo Fisher Scientific), 5 ngmL insulin (Millipore Sigma, Burlington, MA, USA), 40 ngmL hydrocortisone (Millipore Sigma), and 3.5 ngmL human recombinant epidermal growth factor (EGF) (Millipore Sigma).	Health Healthcare		
After washing with PBS, the fluorescence of the cells was analyzed using a flow cytometer (FACSLyric,	FlowJo		Suggestion: FlowJo, RRID:SCR_008520(link)

BD Biosciences, San Jose, CA, USA) and FlowJo software (BD Biosciences).			
Statistical analysis was performed using GraphPad Prism software (GraphPad Software, La Jolla, CA, USA).	GraphPad Prism		Suggestion: GraphPad Prism (RRID:SCR_002798)(link)
	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
For one-time measurements, a one-way ANOVA followed by Tukey test was used.	ANOVA
	Tukey test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq