Document Identifier: **PMC8354104**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: This study was approved by the Animal Ethics Committee of Nantong University of China (Approval No. S20180923-007) in 2018 .
Euthanasia Agents: Lingfeng Chemical Reagent , Shanghai , China) , and 17 mg/kg sodium pentobarbital (MilliporeSigma , Burlington , MA , USA) , the main trunk of the right extracranial facial nerve was exposed.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
Materials and Methods Animals Sixty 6-week-old male Sprague-Dawley rats (specific pathogen-free level), weighing 160–180 g, were included in this study.
Subject Demographics
Age: Materials and Methods Animals Sixty 6-week-old male Sprague-Dawley rats (specific pathogen-free level), weighing 160–180 g, were included in this study.
Weight: Materials and Methods Animals Sixty 6-week-old male Sprague-Dawley rats (specific pathogen-free level), weighing 160–180 g, were included in this study.
Randomization
All included rats were randomly divided into six groups (n = 10) .
Blinding
Video recordings of the blink reflex , vibrissae movement , and tip position were analyzed by two independent blinded observers at the time of acquisition .
Power Analysis
not detected.

Replication
Each sample was replicated in three wells.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Anti-Thy1.1 antibody (mouse , 1:1000 , Cat# M7898 , Sigma-Aldrich , St . Louis , MO , USA) and rabbit complement (25 % , Cat# 31024100 , Invitrogen , Carlsbad , CA , USA) were subsequently applied to remove fibroblasts and purify the isolated cells .	Anti-Thy1.1		
Experimental Models: Organisms/Strains			
Materials and Methods Animals Sixty 6-week-old male Sprague-Dawley rats (specific pathogen-free level), weighing 160–180 g, were included in this study.	Sprague-Dawley		Suggestion: RRID:RGD_70508(link)
Software and Algorithms			
The series test of cluster analysis was performed using Stem software (http://www.cs.cmu.edu/~jernst/stem/) , and the co-expression network analysis was performed using CytoScape software (https://cytoscape.org/) with bioinformatics assistance from Shanghai OE Biotech . Co . , Ltd. (Shanghai , China) .	http://www.cs.cmu.edu/~jernst/stem/		
	CytoScape		Suggestion: Cytoscape (RRID:SCR_003032)(link)
	https://cytoscape.org/		
Then , these candidate genes were uploaded for gene enrichment analysis with the corresponding database (Gene Ontology (GO) enrichment: http://geneontology.org/; Kyoto Encyclopedia of Genes and Genomes (KEGG) enrichment: http://www.genome.jp/kegg/) , and the PubMed database (https://www.ncbi.nlm.nih.gov/gene/) was used to screen genes related to cell proliferation , migration , apoptosis , and the cell cycle.	KEGG		
	PubMed		Suggestion: PubMed (RRID:SCR_004846)(link)
S100β (mouse , 1:1000 , Cat# s2532 , Sigma) , was used to immunostain the purified cell preparation , and the results indicated that the cell preparation consisted of 98 % SCs .	Sigma		
Finally, migrating cells at the wound front were photographed, and the clear areas were measured by ImageJ software at the indicated time points	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)

to determine the percentage relative to that at 0 hours.			
Statistical analysis All statistical analyses were conducted by Student's t-test using GraphPad Prism 9 (GraphPad Software, San Diego, CA, USA).	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)
The DEGs were statistically analyzed using DESeq software (https://www.huber.embl.de/users/anders/DESeq/).	DESeq		Suggestion: DESeq, RRID:SCR_000154(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical analysis All statistical analyses were conducted by Student’s t-test using GraphPad Prism 9 (GraphPad Software, San Diego, CA, USA).	t-test
Oligonucleotides	
Additional Table 1 Lists of primers used in quantitative real-time polymerase chain reaction Gene ID Gene full name Gene symbol Primer sequences (5’-3’)	5’-3’
Rho GTPase activating protein 30 Arhgap30 Forward: GGAGTTCCTTATGCGCCACT Reverse: TCTACACCAGCAGAGAGGGAA	GGAGTTCCTTATGCGCCACT
	TCTACACCAGCAGAGAGGGAA
Aldo-keto reductase family 1 , member B8 Akr1b8 Forward: AGCCCAGGTTCTTATTCGGT Reverse: TCTGCGTCATAGGGAAACTCT	AGCCCAGGTTCTTATTCGGT
	TCTGCGTCATAGGGAAACTCT
Complement C5a receptor 1 C5ar1 Forward: CCAGGACATGGACCCCATAA Reverse: GGCGTTGACAGTACGTTTGG 652957	CCAGGACATGGACCCCATAA
	GGCGTTGACAGTACGTTTGG
Colony stimulating factor 2 receptor alpha subunit Csf2ra Forward: CAACTCGGAGACGCTGAGGAA Reverse: TGTCAGAGATGGAGGATGGGAG 360509 Dedicator of cytokinesis 2 Dock2 Forward: TTCTTCAAGACCATGCTGGC Reverse: ACAGAAGATGCGAGGCTTGT	CAACTCGGAGACGCTGAGGAA
	TGTCAGAGATGGAGGATGGGAG
	TTCTTCAAGACCATGCTGGC
	ACAGAAGATGCGAGGCTTGT
288077 Hematopoietic cell specific Lyn substrate 1 Hcls1 Forward: GGGCCACGAGTATGTTGCT Reverse: CCCTTGGCATAATCTCGCTGA	GGGCCACGAGTATGTTGCT
	CCCTTGGCATAATCTCGCTGA
Inositol polyphosphate-5-phosphatase D Inpp5d Forward: TGTTCAGGCATCCGAAGGTG Reverse: CGGGGGTTCTCTGTTGGAAG	TGTTCAGGCATCCGAAGGTG
	CGGGGGTTCTCTGTTGGAAG
Spi-1 proto-oncogene Spi1 Forward: ATTGGTGGTGATGGAGACAGC Reverse: GTAGGAAACCTGGTGACTGAGG	ATTGGTGGTGATGGAGACAGC
	GTAGGAAACCTGGTGACTGAGG
src-like adaptor Sla Forward: CACCTGCACCTTTGGAGAGA Reverse: AACAGCCAGCCATGATACACT	CACCTGCACCTTTGGAGAGA
	AACAGCCAGCCATGATACACT
Interleukin 6 IL-6 Forward: GTTTCTCTCCGCAAGAGACTT Reverse: ATACTGGTCTGTTGTGGGTGG 24770 C-C motif chemokine ligand 2 MCP-1 Forward: CCACCACTATGCAGGTCTCTG Reverse: GGGGCATTAACTGCATCTGGC Scoring of facial nerve function The function of the facial nerve was estimated and scored	GTTTCTCTCCGCAAGAGACTT
	ATACTGGTCTGTTGTGGGTGG
	CCACCACTATGCAGGTCTCTG
	GGGGCATTAACTGCATCTGGC

on three aspects (blink reflex , vibrissae movement , and tip position) according to a previously reported method (Gao et al. , 2019) .	
The target sequences of siRNA duplexes were as follows: negative-control siRNA , 5#-GGC UCU AGA AAA GCC UAU GC-3#	5#-GGC UCU AGA AAA GCC UAU GC-3#
, Akr1b8-specific siRNA-1 , 5#-CTG GGT ATC GAC ATA TTG A-3# , Akr1b8-specific siRNA-2 , 5#-ACC CTT CAC TAT TAC AGG A-3# , Akr1b8-specific siRNA-3 , 5#-CAG CAC GTA TAC AGG AGA A-3# .	5#-CTG GGT ATC GAC ATA TTG A-3#
	5#-ACC CTT CAC TAT TAC AGG A-3#
	5#-CAG CAC GTA TAC AGG AGA A-3#

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq