Document Identifier: **PMC8355681**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not required.
Inclusion and Exclusion Criteria
not required.
Attrition
not required.
Sex as a biological variable
not detected.
Subject Demographics
Age: not required.
Weight: not required.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
Number: Every experiment was carried out independently three times.
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
HIF-1a , glucose transporter type 1 (GLUT1) , hexokinase 2 (HK2) , pyruvate kinase M2 (PKM2) , lactate dehydrogenase A (LDHA) and b-actin antibodies were from Abcam .	HIF-1a		
	glucose transporter type 1 (GLUT1)		
	, hexokinase 2		
	hexokinase 2 (HK2)		
	HK2		
	pyruvate kinase M2 (PKM2)		
	lactate dehydrogenase		
	b-actin antibodies		
	b-actin		
The following primary antibodies were used (all from Abcam): Caspase-3 and C-Caspase-3 (cat. no. ab13847 1:200) , HIF-1a (cat. no. ab179483 1:500) , GLUT1 (cat. no. ab115730 1:200) , HK2 (cat. no. ab209847 1:600) , PKM2 (cat. no. ab137852 1:500) , LDHA (cat. no. ab101562 1:300) and b-actin (cat. no. ab8226 1:500) .	GLUT1		
	PKM2		
	LDHA		
A rabbit IgG-HRP antibody (Abcam cat. no. ab270144 1:800) was used as the secondary antibody and was incubated at 25C for 1 h . b-actin antibody (cat. no. ab6276 1:1,000 Abcam) was used as an internal reference.	rabbit IgG-HRP		
Experimental Models: Cell Lines			
Chronic myelogenous leukaemia cell lines K562 and K562ADM were from American Type Culture Collection .	K562		Suggestion: RRID:CVCL_0004(link)
	K562ADM		
Recombinant DNA			
The coding sequence of HIF-1a (National Center for Biotechnology Information reference sequence , NM_001530.3) , HIF-1a small interfering RNA (siHIF-1a: 5-CTACTCAGGACACAGATTTAGACTTGGAG-3) and the	pcDNA-3.1		

scramble siRNA (5-CATCCTGAACAGACTTAAGACGTGGTAGT-3) were synthesized and cloned into pcDNA-3.1 vectors for HIF-1a overexpression and knockdown by Sangon Biotech Co . , Ltd. miRNAs that could specifically target HIF-1a were screened using Targetscan 7.2 (http:www.targetscan.orgvert_72) and Miranda 2.0 (http:www.miranda.org)			
The pRL-CMV Renilla luciferase vector (Promega Corporation) was used to normalize cell numbers and transfection efficiency .	pRL-CMV		
Software and Algorithms			
The coding sequence of HIF-1a (National Center for Biotechnology Information reference sequence , NM_001530.3) , HIF-1a small interfering RNA (siHIF-1a: 5-CTACTCAGGACACAGATTTAGACTTGGAG-3) and the scramble siRNA (5-CATCCTGAACAGACTTAAGACGTGGTAGT-3) were synthesized and cloned into pcDNA-3.1 vectors for HIF-1a overexpression and knockdown by Sangon Biotech Co . , Ltd. miRNAs that could specifically target HIF-1a were screened using Targetscan 7.2 (http:www.targetscan.orgvert_72) and Miranda 2.0 (http:www.miranda.org)	Targetscan		Suggestion: TargetScan (RRID:SCR_010845)(link)
K562ADM cells were collected and stained with PI and Annexin V for 15 min at 25C away from light and tested by a BD flow cytometer (BD Biosciences) and analyzed using FlowJo(tm) software version 7.6.5 (FlowJo LLC)	FlowJo		Suggestion: FlowJo, RRID:SCR_008520(link)
Statistical analysis Statistical analyses were performed using SPSS software version 16.0 (SPSS, Inc.).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical significance was evaluated by one-way ANOVA with Turkeys post hoc test.	ANOVA
	Turkeys post hoc test
Oligonucleotides	
The coding sequence of HIF-1a (National Center for Biotechnology Information reference sequence , NM_001530.3) , HIF-1a small interfering RNA (siHIF-1a: 5-CTACTCAGGACACAGATTTAGA CTTGGAG-3) and the scramble siRNA (5-CATCCTGAACAGACTTAAGACG TGGTAGT-3) were synthesized and cloned into pcDNA-3.1 vectors for HIF-1a overexpression and knockdown by Sangon Biotech Co . , Ltd. miRNAs that could specifically target HIF-1a were screened using Targetscan 7.2 (http:www.targetscan.orgvert_72) and Miranda 2.0 (http:www.miranda.org)	5-CTACTCAGGACACAGATTTAGA CTTGGAG-3
The qPCR primer pair for miR-18a-5p was as follows: Forward , GGGGAUAGACGUGAUCUA and reverse , GTGCAGGTCCGAGGT .	GGGGAUAGACGUGAUCUA
	GTGCAGGTCCGAGGT
The primer pair for U6 was as follows: Forward , CTCGCTTCGGCAGCACA and reverse , AACGCTTCACGAATTTGCGT .	CTCGCTTCGGCAGCACA
	AACGCTTCACGAATTTGCGT
The primer pair for HIF-1a was as follows: Forward , GGACAAGTCACCACAGGACA and reverse , GGGAGAAAATCAAGTCGTGC .	GGACAAGTCACCACAGGACA
	GGGAGAAAATCAAGTCGTGC
The primer pair for b-actin was as follows: Forward , CACCTTCTACAATGAGCTGCGTGT G and reverse , ATAGCACAGCCTGGATAGCAACGT AC .	CACCTTCTACAATGAGCTGCGTGT G

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq