Document Identifier: **PMC8357711**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: Individuals were ineligible if they belonged to a group for which aspirin use is not recommended (i.e. if they had glucose-6-phosphate dehydrogenase deficiency or another condition for which low-dose aspirin is contraindicated, were being treated with methotrexate, or were pregnant) were employed by a pharmaceutical company had a direct role in treating patients with CVD or had a cognitive impairment, hearing difficulty, visual impairment, or acute psychopathology that, in the investigators or interviewers judgement, could interfere with the ability to provide online consent and complete an interview or survey.
Inclusion and Exclusion Criteria
Individuals were ineligible if they belonged to a group for which aspirin use is not recommended (i.e. if they had glucose-6-phosphate dehydrogenase deficiency or another condition for which low-dose aspirin is contraindicated, were being treated with methotrexate, or were pregnant) were employed by a pharmaceutical company had a direct role in treating patients with CVD or had a cognitive impairment, hearing difficulty, visual impairment, or acute psychopathology that, in the investigators or interviewers judgement, could interfere with the ability to provide online consent and complete an interview or survey.
Attrition
not detected.
Sex as a biological variable
Individuals were ineligible if they belonged to a group for which aspirin use is not recommended (i.e. if they had glucose-6-phosphate dehydrogenase deficiency or another condition for which low-dose aspirin is contraindicated, were being treated with methotrexate, or were pregnant) were employed by a pharmaceutical company had a direct role in treating patients with CVD or had a cognitive impairment, hearing difficulty, visual impairment, or acute psychopathology that, in the investigators or interviewers judgement, could interfere with the ability to provide online consent and complete an interview or survey.
Subject Demographics
Age: Participants The study enrolled two groups of participants: adults aged 50-75 years eligible to receive low-dose aspirin (100 mgday) for primary prevention of CVD and adults aged >= 18 years eligible to receive low-dose aspirin for secondary prevention of CVD.
Randomization
DCE Thirty-six DCE questions were generated using a D-efficient design, which were divided across three blocks with 12 questions each and randomly allocated to participants.

Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
We reviewed current practice guidelines of the United States Preventative Services Task Force, European Society of Cardiology, World Health Organization, American College of Chest Physicians, and American Heart Association, as well as published literature, which was identified by searching major medical databases for both qualitative and quantitative preference studies on the use of aspirin, antiplatelet drugs, and non-steroidal anti-inflammatory drugs in primary or secondary prevention of CVD or CRC.	American Heart Association		Suggestion: American Heart Association (RRID:SCR_007210)(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
For each class, maximum likelihood estimates reflecting preferences for marginal changes in the attributes were obtained, together with their standard error, and statistical significance of the estimated preferences was determined by two-sided z-test at the 5 level.	z-test
Confidence intervals for the RI scores were calculated by the Krinsky-Robb procedure 32, 33 using 10,000 draws for the simulation, and Wald tests were used to compare RI profiles across latent classes.	Wald tests
Chi-square test of sample proportions was used to detect significant differences in the personal characteristic compositions of the latent classes.	Chi-square test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq