Document Identifier: **PMC8359173**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: 2.2 Ethics review The study was approved by the Institutional Review Boards (IRBs) of Karolinska Institutet, Stockholm, Sweden, Faculty of Medicine, Addis Ababa University, and Ethiopian National Ethics Review Committee and Food Drug and Health Care Administration and Control Authority of Ethiopia.
Consent: All participants gave written informed consent.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
The TB-HIV co-infected patients were also part of a previous open-label randomized clinical trial which investigated efficacy and safety of efavirenz-based ART, initiated at a different time point during the intensive phase of anti-TB therapy (Amogne et al., 2015).
Blinding
not detected.
Power Analysis
not detected.
Replication

not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Statistical analyses were performed using Statistica version 13 (StatSoftInc, Tulsa, OK, USA).	Statistica		Suggestion: STATISTICA (RRID:SCR_014213)(link)
Graph Pad Prism version 8.0 for Windows (Graph Pad, La Jolla, CA, USA) was used for graphical presentations.	Graph Pad Prism		
Efavirenz and lamivudine with zidovudine or stavudine was the HIV treatment and supplied by Mylan Laboratories Limited (Dist-Dhar, M.P, India) 2.9 Nomenclature of targets and ligands Key protein targets and ligands in this article are hyperlinked to corresponding entries in the the IUPHARBPS Guide to PHARMACOLOGY (http:www.guidetopharmacology.org) and are permanently archived in the Concise Guide to PHARMACOLOGY 201920 (Alexander, Cidlowski et al., 2019 Alexander, Fabbro et al., 2019a, 2019b Alexander, Kelly et al., 2019a, 2019b).	http:www.guidetopharmacology.org		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
A chi-square test was used to compare the observed and expected allele frequencies according to the Hardy-Weinberg equilibrium .	chi-square test
Between treatment , group comparison of median plasma cholesterol concentrations , 4b-OHC concentrations , and 4b-OHCChol ratios at baseline and during treatment was done using the Mann-Whitney U test .	Mann-Whitney U test
Therefore , concentration data were transformed into log10 values for statistical analysis and the Shapiro-Wilk test for normality was applied.	Shapiro-Wilk test
Comparison of geometric means of 4b-OHCChol ratio between treatment groups was done using one-way ANOVA followed by Tukeys test .	ANOVA
	Tukeys test
Pairwise comparison of data from baseline within and between treatment groups was made using paired and unpaired t-test , respectively .	t-test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq