Document Identifier: **PMC8359274**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: The study was approved by the local ethics committee (CMO Region Arnhem-Nijmegen, The Netherlands).
Consent: All participants gave written informed consent and were financially compensated for participation. 2.2 ADHD scores The Dutch version of the ADHD DSM-IV Rating Scale was used to measure self-reported current ADHD symptoms (Kooij et al., 2005).
Inclusion and Exclusion Criteria
The exclusion criteria were a history of neurological or psychiatric diseases.
Attrition
Genome-wide genotyping of single nucleotide polymorphisms (SNPs) was performed for the 1,531 unrelated individuals on three genotyping platforms: for 678 participants, genotyping was performed using Affymetrix GeneChip (Affymetrix Inc., Santa Clara, CA), for 634 participants, genotyping was performed using Infinium OmniExpress BeadChip (Illumina Inc., San Diego, CA), and 219 participants were genotyped using Infinium PsychArray BeadChip (Illumina Inc.).
Sex as a biological variable
not detected.
Subject Demographics
Age: Three variables were calculated, as described in (Vrijsen et al., 2018): lifetime-SLEs indicated the total number of experienced life events across the whole life span childhood trauma (CT) reflected the number of traumatic events that participants experienced (aggression, physical andor sexual abuse) within or outside the family before 16 years old recent-SLEs indicated the number of stressful events (e.g., health problems, death of a family member, problems in the romantic relationship, divorce, conflicts in work, financial problems, or legal issues) that participants had within the last year.
Randomization
not detected.
Blinding
not detected.

Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Using the Rapid Imputation Consortium Pipeline (RICOPILI) (Lam et al., 2019), quality control was carried with default parameters and ungenotyped SNPs were imputed using 1,000 genomes Phase I version 2 as a reference and IMPUTE2 software (Howie, Donnelly, & Marchini, 2009).	IMPUTE2		Suggestion: IMPUTE2 (RRID:SCR_013055)(link)
Polygenic risk scores were created using PRSice2 software (Choi & OReilly, 2019).	PRSice2		
Mediation analyses were performed in IBM SPSS Statistics 22, with using the PROCESS macro for SPSS (Hayes, 2013 Sales, 2016).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Bonferroni correction for multiple testing was implemented for the number of comparisons (correlation analysis), resulting in a significance level of p < .	Bonferroni

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq