Document Identifier: **PMC8363827**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
not required.
Inclusion and Exclusion Criteria
Inclusionary criteria for the larger project included maternal age >=18 years old, native English speaking, singleton pregnancy, and normal fetal brain anatomy as assessed by ultrasound and MRI examination.
Attrition
Forty-three percent of children in the N = 115 sample were female and MRI studies occurred when fetuses were between 22- and 39-weeks gestational age (GA).
Sex as a biological variable
2 Materials and methods 2.1 Participants This study included data from 115 pregnant, second-and third-trimester mothers from a larger ongoing project on fetal brain development who had available urine toxicology data (1-5 urine toxicology tests, mean = 1.83).
Subject Demographics
Age: Inclusionary criteria for the larger project included maternal age >=18 years old, native English speaking, singleton pregnancy, and normal fetal brain anatomy as assessed by ultrasound and MRI examination.
Weight: Additional exclusion criteria were applied to neuroimaging data, including exclusion of fetuses scanned before 27 weeks (n = 4) those with health concerns, such as delivery before 33 weeks, birth weight less than 2500 g (n = 18) lack of sufficient low-motion fMRI data (n = 11) or presence of multi-drug or other drugs in the drug toxicology screen (n = 14), resulting in a PCEnon-PCE neuroimaging sample of N = 68.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.

Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Each segment was manually reoriented , realigned to the mean BOLD volume , resampled to 2 mm isotropic voxels , and normalized to a 32-week fetal brain template (Serag et al. , 2012) using Statistical Parametric Mapping (SPM12) (Statistical Parametric Mapping 8 from the Wellcome Trust Centre for Neuroimaging , 2009) software implemented in MATLAB .	Statistical Parametric Mapping		Suggestion: Spatial Statistical Parametric Mapping (RRID:SCR_002592)(link)
	MATLAB		Suggestion: MATLAB, RRID:SCR_001622(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
) 0.651 Rotational drift (degrees) 1.18 (0.4) 1.24 (0.4) 0.502 Note: Chi-square tests compared raceethnicity , marital status , income , education , and child sex between groups .	Chi-square tests
Two-sample t tests compared all other variables .	t tests
2.7 Statistical analyses Demographic and questionnaire data were analyzed using standard Chi-squared test statistics or two-tailed independent samples t-tests	Chi-squared test
Hippocampal RSFC was compared between PCE and non-PCE groups using a two-sample t-test , controlling for gestational age at scan .	t-test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq