Document Identifier: **PMC8369018**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: 15 Patients were stratified according to eGFR as having mild (>=60 mLm), moderate (30-69 mLm), or severe (<30 mLm) CKD.16 Ethical approval and consent to participate All patients provided written informed consent.
IRB: The ethics committee of Nanjing Medical University approved all the experimental protocols.
Inclusion and Exclusion Criteria
Patients with any of the following were excluded: massive hemorrhage within 1 month severe thrombocytopenia (blood platelet count <20x109L) dialysis-dependent end-stage renal failure or allergy to bivalirudin or hirudin.
Attrition
not detected.
Sex as a biological variable
For females with SCr <=(>)0.7, then eGFR=(144)x(SCr0.7)ax(0.993)age, where a=-0.329 (-1.209).
Subject Demographics
Age: Massive hemorrhage sufficient for exclusion was defined as clinically overt bleeding, accompanied by a decrease in hemoglobin >=2 gdL, requiring a transfusion of >=2 U of packed red blood cells, and occurring at a site of concern (intracranial, intraocular, intraspinal, intra-articular, intramuscular with compartment syndrome, pericardial, or retroperitoneal), or resulting in death.11 Finally, the study population consisted of 134 patients, aged 40 to 89 years.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.

Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Data analysis The data analysis was performed using the Statistical Package for Social Sciences software (ver. 16.0 SPSS, Chicago, IL, USA).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Data analyses utilized chi-squared tests to determine differences in sex, smoking status, drinking status, and medical history.	chi-squared tests
Independent samples t-tests, one-way analysis of variance, and paired samples t-tests were applied to normal data, as appropriate.	t-tests
	analysis of variance
Other baseline characteristics (non-normal data) were examined by Mann-Whitney and Wilcoxon rank tests.	Mann-Whitney
	Wilcoxon rank tests

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq