Document Identifier: **PMC8374740**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: The study protocol (Supplement 1) was approved by an independent ethics committee or institutional review board at each study site (eTable in Supplement 2).
Consent: All patients provided written informed consent.
Inclusion and Exclusion Criteria
Patients with a baseline draining fistula count of more than 20; those requiring surgical management before week 13; those requiring excisional surgery with primary closure, partial surgical reduction of excision with surgical suture, or reconstruction techniques; and those with active skin disease or conditions other than HS that could interfere with the assessment of HS were excluded.
Attrition
During the perioperative period (weeks 13 and 14), patients continued adalimumab, 40 mg, weekly or placebo, and the surgeon measured and recorded the surface area of the actual surgery.
Sex as a biological variable
not detected.
Subject Demographics
Age: SHARPS enrolled adults (aged 18-65 years) diagnosed with HS 1 year or more prior to baseline.
Randomization
Methods Study Design The SHARPS study was a global, phase 4, prospective, multicenter, randomized, double-blind, placebo-controlled study of adalimumab in conjunction with surgery in adults with moderate to severe HS who were surgical candidates.
Blinding
not detected.
Power Analysis
Based on the combined response rates for the HiSCR from previous studies, 9 a conservative estimate of 20% treatment difference was used for the power calculation in which case 100 patients per treatment arm would

provide at least 80% power to detect a treatment difference at the alpha level of .05 and to demonstrate the point estimate of the treatment difference of at least 15% for the primary end point.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Analyses were performed using SAS statistical software version 9.4 (SAS Institute).	SAS statistical		
	SAS Institute		Suggestion: Statistical Analysis System, RRID:SCR_008567(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Categorical variables were analyzed using Cochran-Mantel-Haenszel test, stratified by baseline Hurley stage and anatomical location of the HS surgical site and continuous variables using analysis of covariance, with treatment, baseline Hurley stage, anatomical location of the HS surgical site, and baseline values in the model.	Cochran-Mantel-Haenszel test
	analysis of covariance

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq