Document Identifier: **PMC8377279**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: They were admitted to the departments of internal medicine and were included if they understood the Dutch language and provided informed consent.
IRB: Ethical Considerations The study was conducted in accordance with principles of the Declaration of Helsinki and was approved by the local Medical Ethics Committee of Maastricht (METC 2018-0844).
Inclusion and Exclusion Criteria
They were admitted to the departments of internal medicine and were included if they understood the Dutch language and provided informed consent.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: Individuals over 60 years were identified as patients who could benefit from palliative care (SQ).
Randomization
not detected.
Blinding
not detected.
Power Analysis
With 66 patients in each group, it was estimated that the study had a power of 80 to detect a 20 difference in proportion of patients who significantly improved on the USD TSDS, with a minimal clinically important difference of 6, comparing the 2 study groups, given the conservative estimate of the loss-to-follow-up of 10. 23,24 The primary analysis was performed using a Chi-square goodness-of-fit test.
Replication

not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Statistical Analysis The statistical analysis was carried out with SPSS (version 25.0).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Baseline patient characteristics of both study arms were compared using Chi-square tests, students t-tests, or Mann-Whitney U tests, as appropriate.	Chi-square tests
	t-tests
	Mann-Whitney U tests
With 66 patients in each group, it was estimated that the study had a power of 80 to detect a 20 difference in proportion of patients who significantly improved on the USD TSDS, with a minimal clinically important difference of 6, comparing the 2 study groups, given the conservative estimate of the loss-to-follow-up of 10. 23,24 The primary analysis was performed using a Chi-square goodness-of-fit test.	Chi-square goodness-of-fit test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq