Document Identifier: **PMC8381433**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: All patients provided written consent for treatment.
IRB: The Ethics Committee of Iwate Medical University (Iwate, Japan) approved this clinical trial (MH2019-606) prior to patient recruitment on 25 February 2019, and the details of patients were de-identified.
Inclusion and Exclusion Criteria
Twenty-four patients were excluded from the study because they had been followed up for less than 6 months.
Attrition
Twenty-four patients were excluded from the study because they had been followed up for less than 6 months.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
In addition to background characteristics, the EORTC and CUETO recurrence scores and the factors that the scores are based on were calculated for each patient, and the incidence of a second TURBT and BCG therapy, current or past medication history of therapeutic agents for dysuria and recurrence status during the follow-up period were recorded.	EORTC		Suggestion: EORTC (RRID:SCR_004070)(link)
Statistical analysis The data were analyzed using JMP(r) 14 statistical software (SAS Institute Inc., Cary, NC, USA).	SAS Institute		Suggestion: Statistical Analysis System, RRID:SCR_008567(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Differences between the non-recurrence and recurrence groups in terms of continuous variables, such as follow-up period and age, were analyzed using t-tests.	t-tests
The Cox regression test was used for categorical variables.	Cox regression test
The recurrence-free survival (RFS) was calculated using Kaplan-Meier curves, and the significance of differences between survival curves was determined using the log-rank test.	log-rank test
Moreover, the hazard ratio (HR) was calculated using Cox regression analysis. $p<0.05$ was considered to indicate a statistically significant difference.	Cox regression

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq