Document Identifier: **PMC8382122**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: Ethics statement All animal studies and experiments were approved and performed under the Kansas State University (KSU) Institutional Biosafety Committee (IBC, Protocol 1460) and the Institutional Animal Care and Use Committee (IACUC, Protocol 4468) in compliance with the Animal Welfare Act.
Inclusion and Exclusion Criteria
A word size of 40 was used for the BLAST analysis in order to exclude reads where the target mutation fell at the end of a read or reads that partially covered the target sequence.
Attrition
Five of the six deer were pregnant the number of fetuses per deer are indicated in Table 1.
Sex as a biological variable
Virus challenge of animals Six female white-tailed deer (WTD), approximately 2 years of age, were acquired from a Kansas deer farm (Muddy Creek Whitetails, KS) and acclimated for ten days in BSL-3Ag biocontainment with feed and water ad libitum prior to experimental procedures.
Subject Demographics
Age: Virus challenge of animals Six female white-tailed deer (WTD), approximately 2 years of age, were acquired from a Kansas deer farm (Muddy Creek Whitetails, KS) and acclimated for ten days in BSL-3Ag biocontainment with feed and water ad libitum prior to experimental procedures.
Randomization
not detected.
Blinding
Two independent veterinary pathologists (blinded to the treatment groups) examined the slides and morphological descriptions were provided.
Power Analysis
not detected.
Replication

Briefly, heat inactivated (56degC 30 min) serum samples were subjected to 2-fold serial dilutions starting at 1:20 and tested in duplicate.
Number: Briefly, heat inactivated (56degC 30 min) serum samples were subjected to 2-fold serial dilutions starting at 1:20 and tested in duplicate.
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Following three washes with PBS-T, 100 mL of HRP-labelled Rabbit Anti-Deer IgG (H+L) secondary antibody (95058-328, VWR, Batavia, IL, USA) diluted 1:1000 (100ngmL) was added to each well and incubated for 1 hour at RT.	Anti-Deer IgG		
Indirect ELISA was used to detect Bovine Coronavirus (BCoV) antibodies in sera with Spike (S) recombinant viral protein (LSBio, LS-G64076-20, Seattle, WA, USA) using the methods described above.	BCoV		
SARS-CoV-2-specific immunohistochemistry (IHC) IHC was performed as previously described 11 on four-micron sections of formalin-fixed paraffin-embedded tissue mounted on positively charged Superfrost(r) Plus slides and subjected to IHC using a SARS-CoV-2-specific anti-nucleocapsid rabbit polyclonal antibody (3A, developed by our laboratory) with the method previously described 12 .	anti-nucleocapsid		
Experimental Models: Cell Lines			
To determine infectious virus titers of virus stocks and study samples, 10-fold serial dilutions were performed on Vero-E6TMPRSS2 cells.	Vero-E6TMPRSS2		
Selected swab and tissue homogenate samples were tested for viable virus by culture on Vero E6TMPRRS2 cells.	Vero E6TMPRRS2		
Susceptibility of cervid cells to SARS-CoV-2 The SARS-CoV-2 USA-WA12020 strain was passaged 3 times in Vero-E6 cells to establish a stock virus for infection experiments.	Vero-E6		Suggestion: RRID:CVCL_0574(link)
Cell supernatants were titrated on Vero E6 cells to determine TCID50mL.	Vero E6		Suggestion: RRID:CVCL_0574(link)
Software and Algorithms			
Swabs were placed in 2mL of viral transport medium (DMEM, Corning combined with 1 antibiotic-antimycotic, ThermoFisher), vortexed, and aliquoted directly into	ThermoFisher		Suggestion: ThermoFisher Connect Platform (RRID:SCR_023441)(link)

cryovials and RNA stabilizationlysis Buffer RLT (Qiagen, Germantown, MD, USA).			
A word size of 40 was used for the BLAST analysis in order to exclude reads where the target mutation fell at the end of a read or reads that partially covered the target sequence.	BLAST		Suggestion: WU-BLAST (RRID:SCR_011824)(link)

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq