Document Identifier: **PMC8408553**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: Genetically engineered breast cancer murine model All animal experiments were performed in compliance with the Duke Institutional Animal Care and Use Committee (IACUC) protocols and housed in an animal facility at Duke University (Durham , NC , USA) .
Inclusion and Exclusion Criteria
An additional PAMAM-G3 mouse was excluded due to having red lungs of double the usual mass, and histological evaluation indicating evidence of a hemorrhage.
Attrition
not detected.
Sex as a biological variable
Male MMTV-PyMT transgenic mice (stock no: 002374) and female FVBNJ mice (stock no: 001800) were purchased from The Jackson Laboratory .
Subject Demographics
Age: Starting 5 weeks post birth , hemizygous MMTV-PyMT females were treated intraperitoneally twice per week with either PAMAM-G3 (20 mgkg) or saline until mice were euthanized at 12 weeks of age.
Randomization
not detected.
Blinding
With blinding as to treatment status, each metastatic lesion was identified, and then associated surface area determined based on the section where the lesion was largest using ImageJ.
Power Analysis
not detected.
Replication
All data were presented as the mean +- standard error determined from 3-5 technical replicates.

Type: All data were presented as the mean +- standard error determined from 3-5 technical replicates.
Number: All conditions were tested in at least duplicate.
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Cell Lines			
The following cell growth media were used for each cell line: SUM149 (HAMF-12 plus 1 L-glutamine , 5 mgmL insulin , 1 mgmL hydrocortisone , 5 non-heat-inactivated fetal bovine serum FBS) , and MDA-MB-231 (MEM plus 10 non-heat inactivated FBS) .	MDA-MB-231		Suggestion: RRID:CVCL_0062(link)
These HEK293 cells have been engineered to stably co-express a TLR gene and an NF-kB-inducible SEAP (secreted embryonic alkaline phosphatase) reporter gene that can be monitored using SEAP detection media .	HEK293		Suggestion: RRID:CVCL_0045(link)
After aspiration of serum-free media from the equilibrated invasion chambers , SUM149 cells were plated into the top chamber at a density of 35,000 cellswell (500 mL final volume) in the presence of serum-free media alone or various combinations of agonists andor PAMAM-G3 (25 or 50 mgmL) .	SUM149		Suggestion: RRID:CVCL_3422(link)
Experimental Models: Organisms/Strains			
Male MMTV-PyMT transgenic mice (stock no: 002374) and female FVBNJ mice (stock no: 001800) were purchased from The Jackson Laboratory .	FVBNJ		Suggestion: RRID:MGI:6143723(link)
A breeding colony was established and maintained by crossing male transgenics to congenic FVB female mice.	FVB		Suggestion: RRID:IMSR_RJ:FVB(link)
Starting 5 weeks post birth , hemizygous MMTV-PyMT females were treated intraperitoneally twice per week with either PAMAM-G3 (20 mgkg) or saline until mice were euthanized at 12 weeks of age.	MMTV-PyMT		
Software and Algorithms			
Materials and methods Reagents PAMAM-G3 solution was purchased from Sigma Aldrich and used in the in vitro experiments.	Sigma Aldrich		Suggestion: Sigma-Aldrich (RRID:SCR_008988)(link)
The cells in the images were then counted using an ImageJ	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)

algorithm unique for each cell line (https:imagej.netParticle_Analysis) .			
All data were plotted and statistically analyzed using GraphPad Prism (version 6.0 or 8.3) or JMP software.	GraphPad Prism		Suggestion: GraphPad Prism (RRID:SCR_002798)(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Quantification and statistical analysis For all assays and experiments, statistical comparison between treatment groups was first tested for normality via the Shapiro-Wilk test.	Shapiro-Wilk test
If normal, the groups were then compared via a two-tailed Students t test or an ANOVA with an appropriate post hoc test.	t test
	ANOVA
If normal, data were analyzed by a two-tailed Students t test or an ANOVA with a Holm-Sidak or Tukeys test.	Holm-Sidak
	Tukeys test
If not normal, the data were then tested with the non-parametric Fishers exact test.	Fishers exact test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq