Document Identifier: **PMC8414439**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IACUC: Methods Ethics statement This study was approved by the Ethics Committee and Animal Care and Use committee of our hospital.
Consent: All participating patients signed informed consent documents.
Euthanasia Agents: Histomorphometric analysis Eight weeks after DMM surgery, mice were euthanized using 3 sodium pentobarbital.
Inclusion and Exclusion Criteria
Patients who met the following conditions were excluded: allergic constitution pregnant or lactating female severe abnormalities affecting other organs, such as heart, brain, liver, or kidney diagnosis of other arthritic diseases, such as rheumatoid arthritis, gouty arthritis, meniscus injury, synovitis, or ligament injury and medicated with non-steroidal anti-inflammatory drugs (NSAIDs) in the past three months, or cortisol or hyaluronic acid preparations in the past six months.
Attrition
not detected.
Sex as a biological variable
The 95 patients with OA included 37 males and 58 females (mean age: 58.64 years (standard deviation (SD) 5.32) mean BMI: 25.04 kgm2 (SD 4.07)), among whom 39 were diagnosed with diabetes.
Subject Demographics
Age: The 95 patients with OA included 37 males and 58 females (mean age: 58.64 years (standard deviation (SD) 5.32) mean BMI: 25.04 kgm2 (SD 4.07)), among whom 39 were diagnosed with diabetes.
Weight: The 95 patients with OA included 37 males and 58 females (mean age: 58.64 years (standard deviation (SD) 5.32) mean BMI: 25.04 kgm2 (SD 4.07)), among whom 39 were diagnosed with diabetes.
Randomization
not detected.
Blinding
not detected.

Power Analysis
not detected.
Replication
Each test took over ten seconds to complete and was repeated 12 times.
Number: Each test took over ten seconds to complete and was repeated 12 times.
Cell Line Authentication
not detected.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
After washing with hybridization solution, the sections were dripped with mouse anti-digoxigenin biotin-labelled antibody working solution, and incubated in a humid chamber at 37degC for 45 minutes.	anti-digoxigenin biotin-labelled		
The sections were incubated with the primary antibodies anti-F480 (ab6640) and anti-doublecortin (ab77450 both Abcam, UK), followed by incubation with corresponding isotype-specific secondary antibodies and labelling with either AlexaFluor-488 or AlexaFluor-633 (Invitrogen, Thermo Fisher Scientific, USA).	anti-F480		Suggestion: (GenWay Biotech Inc. Cat# GWB-BC480F, RRID:AB_10293382)(link)
	anti-doublecortin		
Thereafter , membranes were incubated with diluted primary antibodies , as follows: anti-b-actin (ab179467 , 1:5,000 Abcam) , anti-TGFa (ab92486 , 1:500	anti-b-actin (ab179467		Suggestion: (US Biological Cat# F4510-81C, RRID:AB_2106720)(link)
	anti-TGFa		Suggestion: (LSBio (LifeSpan) Cat# LS-B7423-50, RRID:AB_11233091)(link)
	ab92486		
Experimental Models: Cell Lines			
The correctly sequenced luciferase reporter plasmids (WT and MUT, respectively) were cotransfected into HEK293T cells, with miR-183 mimic or NC mimic, respectively.	HEK293T		Suggestion: RRID:CVCL_0063(link)
Experimental Models: Organisms/Strains			
Surgical destabilization of the medial meniscus to generate a mouse OA model A total of 105 male wild-type (WT) C57BL6 mice (mean age: 8 weeks (SD 1)) were housed in a controlled environment with food and water provided ad libitum and maintained under a 12-hour lightdark cycle.	C57BL6		Suggestion: RRID:IMSR_JAX:000664(link)
Recombinant DNA			
Based on the WT template, mutated (MUT) TGFa 3UTR was generated and inserted into the empty pMIR-reporter between the same cloning sites.	pMIR-reporter		

Software and Algorithms			
Micrographic images were exported to Adobe Photoshop CS 5.1 (Adobe, USA) and adjusted for brightness and contrast to reflect the true colours.	Adobe Photoshop		Suggestion: Adobe Photoshop, RRID:SCR_014199(link)
Dual luciferase reporter assay Target genes of miR-183 were predicted using the bioinformatics tool, TargetScan.	TargetScan		
Protein levels were quantified using ImageJ software (National Institutes of Health (NIH), USA).	ImageJ		Suggestion: ImageJ, RRID:SCR_003070(link)
Statistical analysis All data were analyzed using SPSS v21.0 software (IBM, USA).	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Comparisons between paired groups with normally distributed data and homogeneous variances were conducted by paired t-test.	t-test
Comparisons among multiple groups were conducted by one-way analysis of variance (ANOVA), followed by Tukeys post hoc test.	analysis of variance
	ANOVA
	Tukeys post hoc test
Comparisons among groups at different timepoints were conducted by repeated measures analysis of variance (ANOVA), followed by Bonferronis post hoc test.	Bonferronis post hoc test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq