Document Identifier: **PMC8421144**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Protocols were also approved and performed according to the guidelines of the Ethics Committee for Animal Testing of the University of Navarra (CEEA-06216) and Navarra Government (GN 2016325511) .
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
C57BL6J 8-week-old male mice (Harlan, Indianapolis, IN, USA) were injected i.v. with control (AAV-Ren) and liver expressing shSLU7 (AAV-shSLU7) adenoassociated virus (1 x 1011 pfu) constructed, produced and purified as described (16).
Subject Demographics
Age: C57BL6J 8-week-old male mice (Harlan, Indianapolis, IN, USA) were injected i.v. with control (AAV-Ren) and liver expressing shSLU7 (AAV-shSLU7) adenoassociated virus (1 x 1011 pfu) constructed, produced and purified as described (16).
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
Experiments were performed 15 days after AAV injection with at least 5 animals per group and were repeated at least twice.

Number: Experiments were performed 15 days after AAV injection with at least 5 animals per group and were repeated at least twice.
Cell Line Authentication
not detected.
Code Information
Availability: Sequence of siRNAs will be provided upon request.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Antibodies used were SLU7 (Novus , Biologicals , Centennial , CO , USA NBP2-20403 1:1000)	SLU7		
Experimental Models: Cell Lines			
The HCT116 DNMT1 -- cell line was kindly provided by Dr Rob M. Ewing (Southampton, UK) and described in (22).	HCT116 DNMT1		
Human HCC cell lines PLCPRF5 and HepG2, human cervical carcinoma cell line HeLa, human melanoma cell line A375 and human embryonic kidney immortalized cell line HEK293T were grown in DMEM (Gibco-Life Technology, Madrid, Spain) supplemented with 10 FBS, glutamine and antibiotics.	HCC		
	PLCPRF5		
	HepG2		Suggestion: RRID:CVCL_0027(link)
	HeLa		Suggestion: RRID:CVCL_0030(link)
	A375		Suggestion: RRID:CVCL_0132(link)
	HEK293T		Suggestion: RRID:CVCL_0063(link)
Human non-small cell lung cancer cell line H358 and human adenocarcinoma cell line A549 were grown in RPMI (Gibco-Life Technology) supplemented with 10 FBS, glutamine and antibiotics.	H358		Suggestion: RRID:CVCL_1559(link)
	A549		Suggestion: RRID:CVCL_0023(link)
Human colon cancer cell line HCT116 and HCT116 DNMT1 -- were grown in McCoys 5A (Gibco-Life Technology) supplemented with 10 FBS, glutamine and antibiotics.	HCT116		Suggestion: RRID:CVCL_0291(link)
Human breast cancer cell line MCF7 was grown in MEM (Gibco-Life Technology) supplemented with 10 FBS, insulin and antibiotics.	MCF7		Suggestion: RRID:CVCL_0031(link)
Human glioblastoma cell line U87 was grown in DMEMF12 (Gibco-Life Technology) supplemented with 10 FBS and antibiotics.	U87		
, Thermo Fisher Scientific , Waltham , MA , USA) , and specific methylated (M) and unmethylated (U) primers for the targeted regions of TRIM47 , TFF3 , DLG3 , RASSF1A and GNMT .	RASSF1A		
Recombinant DNA			

pcDNA6.1A-SLU7 plasmid was from GenScript BioTech (Piscataway, NJ, USA) and pcDNA3.1-HDAC1-Flag was kindly provided by Dr Donald E. Ayer (University of Utah, USA)	pcDNA6.1A-SLU7		
	pcDNA3.1-HDAC1-Flag		
Software and Algorithms			
Where indicated, cells were treated before harvesting with the proteasome inhibitor 10 mM MG-132 (Calbiochem, CA, USA, 474790) for the last 20 h or with 20 mM MG-132 for the last 12 h with the protein synthesis inhibitor 10 mml cycloheximide (Sigma-Aldrich, MO, USA, C-7698) for 8 h and 12 h with cell cycle drugs 20 mM lovastatin (Sigma-Aldrich PHR1285) or 2 mgml aphidicolin (Sigma-Aldrich A078) for 60 h or with 330 nM nocodazole (Sigma-Aldrich M1404) for 12 h or 60 h with the demethylating agent 10 mM 5-aza-2-deoxycytidine (Sigma, St. Louis, MO, USA) during 7 days.	Sigma		Suggestion: sigma-tau (RRID:SCR_000488)(link)
The resulting regions were then annotated using the HOMER software package (25) .	HOMER		
Statistical analysis Statistical analysis was performed using the GraphPad Prism software (version 8).	GraphPad Prism		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Chi-squared test was then applied.	Chi-squared test
Statistically significant differences were estimated by applying two-sided unpaired Students t-test or Mann-Whitney U-test according to sample distribution.	t-test
	Mann-Whitney U-test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq