Document Identifier: **PMC8423660**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: The study protocol was approved by the Ethics Committee of Sakurabashi Watanabe Hospital (Reference no. 17-32).
Consent: One of the investigators obtained informed consent from each patient before they were included in the study.
Inclusion and Exclusion Criteria
Patients were excluded if: (1) primary PCI for an occluded or stenosed culprit vessel was not attempted during the acute phase of MI; (2) they had an arrhythmia disturbing GLS and TMAD measurement, such as atrial fibrillation and multiple extra beats; (3) they underwent incomplete echocardiographic study or their images were not stored; and (4) they had poor image quality that rendered the images unsuitable for analysis.
Attrition
Of these 422 patients, 290 patients who underwent echocardiography 2 weeks after the onset of MI using an iE33 (Philips Healthcare, Andover MA, USA) were enrolled in the present study.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
Data Reproducibility Intra- and interobserver agreements in TMAD4ch, TMAD2ch, and GLS were evaluated by measuring these parameters in 10 randomly selected cases twice by the same observer or by 2 independent observers, respectively.
Blinding
28 Information on subsequent clinical events up to 5 years was gathered by investigators by visiting outpatient clinics or by verbal or written blinded review of the circumstances surrounding each death.
Power Analysis

not detected.
Replication
All parameters were measured in duplicate and averaged.
Number: All parameters were measured in duplicate and averaged.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
The displacement of the mid-portion of the mitral annulus was determined and indexed by LV length as TMAD.	TMAD		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Comparisons were made using 1-way analysis of variance (ANOVA) for continuous variables, and the significance of differences was calculated using Tukey's honestly significant difference (HSD) test for factor analysis.	analysis of variance
	ANOVA
	Tukey's honestly significant difference (HSD) test
Categorical variables were compared using Fisher's exact test.	Fisher's exact test
Event-free survival was analyzed using the Kaplan-Meier method with the log-rank test for group comparisons.	log-rank test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq