Document Identifier: **PMC8448216**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
Consent: All patients underwent detailed history and clinical evaluation after obtaining informed consent.
Inclusion and Exclusion Criteria
Patients with multinodular goitre, toxic nodular goitre, drug-induced thyroiditis, antithyroid drugs use, (3-blockers use and radioiodine therapy), radiation exposure to neck, and prior thyroid surgery were excluded.
Attrition
not detected.
Sex as a biological variable
Similarly, pregnant and lactating females were not included.
Subject Demographics
Age: A total of 111 newly diagnosed thyrotoxic patients without antithyroid medication or (3-blockers use and 45 years of age and sex-matched controls were enrolled for this study.
Randomization
not detected.
Blinding
Radiological Evaluation All patients underwent detailed ultrasonography evaluation of thyroid gland by a single experienced radiologist who was blinded for initial diagnosis of the patients .
Power Analysis
not detected.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
The data were analyzed using SPSS 24 software.	SPSS		Suggestion: SPSS, RRID:SCR_002865(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Normality of data was checked using Shapiro-Wilk test.	Shapiro-Wilk test
Pearsons correlation coefficient was used to analyze correlation between different parameters.	Pearsons correlation
Mann-Whitney U -test and independent t -tests were performed to compare means between two groups as required.	Mann-Whitney U -test
	t -tests
Similarly, analysis of variance (ANOVA) test and Kruskal-Wallis test was used for comparison of means between three or more groups depending on normality of data.	analysis of variance
	ANOVA
	Kruskal-Wallis test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq