Document Identifier: **PMC8453748**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Trials 843 (ClinicalTrials.gov: NCT02577003) and 849 (ClinicalTrials.gov: NCT02564211) were conducted at 25 and 12 trial centres (Supplemental Appendix) in Japan between November 2015 and November 2016, and between October 2015 and March 2017, respectively, in accordance with the principles of Good Clinical Practice and were approved by the appropriate institutional review boards and regulatory agencies.
Consent: Written informed consent was obtained from all study patients.
Inclusion and Exclusion Criteria
Patients were excluded from the trials if they had type 1 diabetes or a history of ketoacidosis, unstable diabetic retinopathy, poorly controlled hypertension (systolic blood pressure ≥160 mm Hg or diastolic blood pressure ≥100 mm Hg), significant cardiovascular disease, active liver disease, renal disease or urological disorders, a history of malignancy or haematological disorders; if they had been treated with insulin or thiazolidinediones within 12 weeks prior to screening or with SGLT2 inhibitors at any time; or if they required treatment with systemic steroids.
Attrition
not detected.
Sex as a biological variable
For both trials, at screening, eligible patients were male or female, aged ≥20 years, with T2D being treated with diet and exercise therapy and meeting either of the following additional criteria: on a stable dose of sitagliptin 50 mg once daily and any single additional OHA or low-dose dual combination therapy prior to screening, with HbA1c ≥6.5% and ≤9.0% (Group A, which expands the sitagliptin monotherapy population after discontinuation of the additional OHA[s] and the subsequent medication stabilization period); or on a stable dose of sitagliptin 50 mg once daily and not on any additional OHAs prior to screening, with HbA1c ≥7.0% and ≤10.0%, and, for Trial 849, FPG ≤230 mg/dL (Group B).
Subject Demographics
Age: For both trials, at screening, eligible patients were male or female, aged ≥20 years, with T2D being treated with diet and exercise therapy and meeting either of the following additional criteria: on a stable dose of sitagliptin 50 mg once daily and any single additional OHA or low-dose dual combination therapy prior to screening, with HbA1c ≥6.5% and ≤9.0% (Group A, which expands the sitagliptin monotherapy population after discontinuation of the additional OHA[s] and the subsequent medication stabilization period); or on a

stable dose of sitagliptin 50 mg once daily and not on any additional OHAs prior to screening, with HbA1c ≥7.0% and ≤10.0%, and, for Trial 849, FPG ≤230 mg/dL (Group B).
Randomization
2 METHODS 2.1 Study designs Trial 843 was randomized, placebo-controlled and double-blind (Supplemental Figure S2A).
Blinding
not detected.
Power Analysis
With a sample size of 69 patients per arm, Trial 843 was estimated to have 90% power to detect a true treatment difference of 0.5% in change from baseline in HbA1c at Week 24, assuming a true standard deviation (SD) of 0.85% (α = 0.05, two-sided) and a discontinuation rate of 10%.
Replication
not required.
Protocol Information

Identifiers: Trials 843 (ClinicalTrials.gov: NCT02577003) and 849 (ClinicalTrials.gov: NCT02564211) were conducted at 25 and 12 trial centres (Supplemental Appendix) in Japan between November 2015 and November 2016, and between October 2015 and March 2017, respectively, in accordance with the principles of Good Clinical Practice and were approved by the appropriate institutional review boards and regulatory agencies.	NCT02577003 NCT02564211

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
In Trial 849, the number and percentage of patients achieving HbA1c <7.0% at various times during the study, and the corresponding 95% CI (based on the method of Clopper and Pearson 17), were calculated.	Pearson 17

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq