Document Identifier: **PMC8457888**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: Owing to the freely accessible resource TCGA database, the study let off the institutional review board approval.
Inclusion and Exclusion Criteria
not detected.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
To ensure the predictive power of the prognostic model , we plotted the calibration graphs by the riskRegression package.
Replication
not required.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
Identification of Ferroptosis-Associated lncRNAs Human GTF explanation files were downloaded from Ensembl (http:asia.ensembl.org), a total of 19,604 mRNAs and 14,086 lncRNAs were distinguished and extracted by manipulating Strawberry Perl software.	Ensembl		Suggestion: Ensembl (RRID:SCR_002344)(link)
To investigate the relationship between the falncRNA and ferroptosis-associated mRNA (famRNA) , it was extremely important for us to construct the falncRNA-famRNA coexpression network by Cytoscape software (version 3.6.0 , https:cytoscape.org) , which was visualized as the Sankey diagram by the R ggplot2 and ggalluvial packages .	Cytoscape		Suggestion: Cytoscape (RRID:SCR_003032)(link)
	https:cytoscape.org		
The results of the clinical role of the prognostic model were generated by the R survivalROC , limma , ggpubr , and ComplexHeatmap packages .	ComplexHeatmap		Suggestion: ComplexHeatmap (RRID:SCR_017270)(link)
We performed gene set enrichment analysis (GSEA) between high-risk and low-risk groups based on the GO (v7.4) and Hallmark (v7.4) gene set collections by the R clusterProfiler package.	clusterProfiler		Suggestion: clusterProfiler (RRID:SCR_016884)(link)
These operations were done by the R ggplot2 , ggtext	ggplot2		Suggestion: ggplot2 (RRID:SCR_014601)(link)
Efficacy of Immunotherapy in Prognostic Signature To assess the efficacy of immunotherapy , we compared whether the expression of immune checkpoints differentiated between the high-risk and low-risk groups , which were used to draw violin plots by R limma and ggpubr packages , including PD-L1 (programmed cell death protein Ligand-1) , PD-1 (programmed cell death 1) ,	limma		Suggestion: LIMMA (RRID:SCR_010943)(link)
The analysis processes were realized by R packages pRRophetic and ggplot2, and IC50 was estimated by the Genomics of Drug Sensitivity in Cancer (GDSC, https:www.cancerrxgene.org).41	https:www.cancerrxgene.org		

Statistical Analysis Univariate, LASSO, and multivariate regression analyses were implemented to qualitatively investigate the appropriate molecules and variables.			
All statistical analyses and visualization of results were performed with R software (version 4.0.3, http:www.r-project.org).	http:www.r-project.org		

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Multivariate Cox regression analysis was utilized to calculate the correlation coefficient and construct a prognostic hazard model and draw a forest plot .	Cox regression
The relationship between the prognostic hazard model and clinical variables was measured by the Wilcoxon rank-sum test , and the results were visualized as a clinical relevance heatmap .	Wilcoxon rank-sum test
36 Condition of Tumor-Infiltrating Immune Cells in Prognostic Signature The file of tumor-infiltrating immune cells was downloaded in TIMER2.0 (http:timer . cistrome.org).37 Spearman correlation analysis was implemented between the prognostic signature and tumor-infiltrating immune cells to calculate the correlation coefficient , and the results were visualized as bubble charts using different software including XCELL , QUANTISEQ , MCPCOUNTER , EPIC , CIBERSORT-ABS , and CIBERSORT.38 The Wilcoxon rank-sum test was implemented to determine which immune cells were distinction in the high-risk and low-risk groups , and the results were presented in box graphs .	Spearman correlation
The survival curve was plotted using the Kaplan-Meier method, and the Log rank test was implemented for differences between groups.	Log rank test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq