Document Identifier: **PMC8474176**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: 2 Materials and Methods The study received Human Ethics Committee and the Institutional Regulatory Board approval.
Consent: Signed informed consent was obtained from all subjects or their nearest kin before the study.
Inclusion and Exclusion Criteria
All included children were not having any visible gross brain abnormalities or comorbidity.
Attrition
not detected.
Sex as a biological variable
not detected.
Subject Demographics
Age: not detected.
Weight: not detected.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.
Replication
HIV diagnosis was performed at the government established centers using a standard protocol including an enzyme-linked-immunosorbent-assay (ELISA) test followed by validation from duplicate HIV rapid tests of greater sensitivity and specificity (Yadav et al., 2017)

Number: HIV diagnosis was performed at the government established centers using a standard protocol including an enzyme-linked-immunosorbent-assay (ELISA) test followed by validation from duplicate HIV rapid tests of greater sensitivity and specificity (Yadav et al., 2017)

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms			
2.3 Diffusion tensor imaging data processing DTI images were preprocessed using FMRIB Software Library (	FMRIB		Suggestion: FSL, RRID:SCR_002823(link)
(FSL version 6.0. 0., http:www.fmrib.ox.ac. ukfsl)) package.	FSL version		Suggestion: FSL, RRID:SCR_002823(link)
In addition, the FSL software also performs outlier detection to identify slices where signal is lost as a consequence of subject movement during the diffusion encoding and replaces those slices by non-parametric predictions by the Gaussian process.	FSL software		Suggestion: FSL, RRID:SCR_002823(link)
Normalization and smoothing of FA and MD maps were performed by statistical parametric mapping (SPM) package 12 (Wellcome Department of Cognitive Neurology, UK) and MATLAB (MathWorks, Natick, MA), these normalized and smoothed maps (FA and MD) were used for the voxel-based analysis.	MATLAB		Suggestion: MATLAB, RRID:SCR_001622(link)
In short, non-gradient diffusion-weighted images of each participant were normalized to the standard (Montreal Neurological Institute) space using earlier defined distribution probability of gray, white, and cerebrospinal fluid tissue types maps supplied with SPM12 package (Ashburner and Friston, 2005), and the obtained normalization parameters were applied on FA and MD maps followed by 8 mm Gaussian filter full-width-at-half-maximum smoothing.	SPM12		

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq