Document Identifier: **PMC8475051**

SciScore Report

Below you will find your SciScore report containing three tables. Your score is calculated based on adherence to scientific rigor criteria (Table 1) and identification of key biological resources (Table 2). Table 3 contains statistical tests and oligonucleotides but is not scored. If SciScore makes any mistakes, please contact us to help us learn and improve.

Table 1: Rigor Adherence Table

Ethics
IRB: The FLAG study was approved by the Mercy Health Research Ethics Committee (Ethics Approval Number R1412) and written informed consent was obtained from all participants.
Consent: The FLAG study was approved by the Mercy Health Research Ethics Committee (Ethics Approval Number R1412) and written informed consent was obtained from all participants.
Field Sample Permit: Whole blood was collected in ethylenediaminetetraacetic acid (EDTA) tubes and centrifuged within 1 hour of collection.
Inclusion and Exclusion Criteria
English-speaking women aged over 18 years, with a singleton pregnancy and normal mid-trimester fetal morphology examination were eligible to participate.
Attrition
These 233 participants were selected from an overall cohort of 518 participants and included 176 control women and 57 women who developed preeclampsia.
Sex as a biological variable
It involved the prospective recruitment of pregnant women from whom we obtained 2000 blood samples at 28 (27+0-29+0 days) and 36 (35+0-37+0) weeks gestation.
Subject Demographics
Age: English-speaking women aged over 18 years, with a singleton pregnancy and normal mid-trimester fetal morphology examination were eligible to participate.
Randomization
not detected.
Blinding
not detected.
Power Analysis
not detected.

Replication
not required.
Data Information
Availability: Methods Data, Materials, and Code Disclosure The data that support the findings of this study are available from the corresponding author upon reasonable request.

Table 2: Key Resources Table

Your Sentences	REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies			
Following endogenous peroxidase quenching and blocking of non-specific binding , sections were incubated at 37 degC for 1 hour with 1:500 GDF15 Monoclonal Antibody (Sapphire Biosciencei , NSW , Australia) in blocking buffer (DAKO) .	GDF15		
For isotype controls , primary antibody was substituted with mouse IgG .	mouse IgG		
Software and Algorithms			
Statistical analyses were performed using GraphPad Prism version 9 (GraphPad Software Inc .	GraphPad Prism		
	GraphPad		Suggestion: GraphPad Prism, RRID:SCR_002798(link)

Other Entities Detected

Your Sentences	Recognized Entity
Statistical Tests	
Statistical Analysis Maternal characteristics and birth outcome data were compared for all women with preeclampsia against controls using a Mann-Whitney U test for continuous data and Fishers exact test for categorical data .	Mann-Whitney U test
	Fishers exact test

SciScore is an automated tool that is designed to assist expert reviewers by finding and presenting formulaic information scattered throughout a paper in a standard, easy to digest format. ***SciScore is not a substitute for expert review.*** SciScore also checks for the presence and correctness of several unique identifiers, including RRIDs (research resource identifiers) in the manuscript, detects sentences that appear to be missing RRIDs, and can even suggest RRIDs under certain circumstances. **All RRID suggestions should be verified;** only the author can know whether the suggestions are correct.

For a full description of scored criteria and tips for improving your score, please see https://www.scicrunch.com/sciscorereport-faq